

B18

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
9 June 2005 (09.06.2005)

PCT

(10) International Publication Number
WO 2005/051931 A2

(51) International Patent Classification⁷: C07D 261/04

Vasant Kunj, New Delhi, Delhi 110070 (IN). DASTIDAR, Sunanda, Ghose [IN/IN]; B-138, Sarita Vihar, New Delhi, Delhi 110044 (IN).

(21) International Application Number:
PCT/IB2004/003893

(74) Common Representative: RANBAXY LABORATORIES LIMITED; c/o DESHMUKH, Jay R., 600 College Road East, Suite 2100, Princeton, NJ 08540 (US).

(22) International Filing Date:
26 November 2004 (26.11.2004)

(25) Filing Language: English

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(26) Publication Language: English

(30) Priority Data:
60/525,347 26 November 2003 (26.11.2003) US

(71) Applicant (for all designated States except US): RANBAXY LABORATORIES LIMITED [IN/IN]; 19, Nehru Place, New Delhi, Delhi 110019 (IN).

(72) Inventors; and

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(75) Inventors/Applicants (for US only): PALLE, Venkata, P. [IN/IN]; D-011, Oakwood Estate, Akashneem Marg, DLF - Phase - II, Gurgaon, Haryana 122001 (IN). BALACHANDRAN, Sarala [IN/IN]; A-6, National Institute of Immunology, Aruna Asif Ali Marg, New Delhi, Delhi 110067 (IN). GUPTA, Nidhi [IN/IN]; J-160, Vikaspuri, New Delhi, Delhi 110018 (IN). KUKREJA, Gagan [IN/IN]; LD-166, Pitampura, Delhi, Delhi 110088 (IN). KHERA, Manoj, Kumar [IN/IN]; #2682, Inside Kot, Ganaur City, Sonapat, Haryana 131101 (IN). BAREGAMA, Lalit, Kumar [IN/IN]; Chyawan-Bhavan, Adarsh Nagar, Kapasan, Chittorgarh, Rajasthan 312202 (IN). MANDADAPU, Raghuramalah [IN/IN]; Mandepudi, Amrawati, Guntur, Andhra Pradesh 522018 (IN). RAY, Abhijit [IN/IN]; Sector C-1, Flat No. 1408,

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PHOSPHODIESTERASE INHIBITORS

(57) Abstract: The present invention relates to isoxazoline derivatives and their analogues, which can be used as phosphodiesterase (PDE) type IV selective inhibitors. Compounds disclosed herein can be useful in the treatment of AIDS, asthma, arthritis, bronchitis, chronic obstructive pulmonary disease (COPD), psoriasis, allergic rhinitis, shock, atopic dermatitis, Crohn's disease, adult respiratory distress syndrome (ARDS), eosinophilic granuloma, allergic conjunctivitis, osteoarthritis, ulcerative colitis and other inflammatory diseases especially in humans. Processes for the preparation of disclosed compounds, pharmaceutical compositions containing the disclosed compounds, and their use as PDE type IV selective inhibitors, are provided.

WO 2005/051931 A2

WO 2005/051931

PCT/IB2004/003893

1

PHOSPHODIESTERASE INHIBITORS

Field of the Invention

The present invention relates to isoxazoline derivatives and their analogues, which
5 can be used as phosphodiesterase (PDE) type IV selective inhibitors. Compounds
disclosed herein can be useful in the treatment of AIDS, asthma, arthritis, bronchitis,
chronic obstructive pulmonary disease (COPD), psoriasis, allergic rhinitis, shock, atopic
dermatitis, Crohn's disease, adult respiratory distress syndrome (ARDS), eosinophilic
granuloma, allergic conjunctivitis, osteoarthritis, ulcerative colitis and other inflammatory
10 diseases especially in humans. Processes for the preparation of disclosed compounds,
pharmaceutical compositions containing the disclosed compounds, and their use as PDE
type IV selective inhibitors, are provided.

Background of the Invention

It is known that cyclic adenosine-3',5'-monophosphate (cAMP) exhibits an
15 important role of acting as an intracellular secondary messenger (E.W. Sutherland, and
T.W. Roll, Pharmacol.Rev,1960,12, 265). Its intracellular hydrolysis to adenosine 5'-
monophosphate (AMP) causes number of inflammatory conditions which are not limited
to psoriasis, allergic rhinitis, shock, atopic dermatitis, Crohn's disease, adult respiratory
distress syndrome (ARDS), eosinophilic granuloma, allergic conjunctivitis, osteoarthritis,
20 ulcerative colitis. The most important role in the control of cAMP (as well as of cGMP)
level is played by cyclic nucleotide phosphodiesterases (PDE) which represents a
biochemically and functionally, highly variable superfamily of the enzyme; eight distinct
families with more than 15 gene products are currently recognized. Although PDE I, PDE
II, PDE III, PDE IV, and PDE VII all use cAMP as a substrate, only PDE IV and PDE VII
25 are highly selective for hydrolysis of cAMP. Inhibitors of PDE, particularly the PDE IV
inhibitors, such as rolipram or Ro-1724 are therefore known as cAMP-enhancers. Immune
cells contain type IV and type III PDE, the PDE IV type being prevalent in human
mononuclear cells. Thus the inhibition of phosphodiesterase type IV has been a target for
modulation and, accordingly, for therapeutic intervention in a range of disease processes.

WO 2005/051931

PCT/IB2004/003893

2

The initial observation that xanthine derivatives, theophylline and caffeine inhibit the hydrolysis of cAMP led to the discovery of the required hydrolytic activity in the cyclic nucleotide phosphodiesterase (PDE) enzymes. More recently, distinct classes of PDE have been recognized (J.A. Bervo and D.H. Reifsnnyder, TIPS, 1990, 11, 150), and their selective inhibition has led to improved drug therapy (C.D. Nicholus, R.A. Challiss and M. Shahid, TIPS, 1991, 12, 19). Thus it was recognized that inhibition of PDE IV could lead to inhibition of inflammatory mediator release (M.W. Verghese et. al, J. Mol. Cell. Cardiol., 1989, 12 (Suppl.II), S 61) and airway smooth muscle relaxation.

U.S. Patent No. 5,686,434 (National stage of WO 95/14680) discloses 3-aryl-2-isoxazolines as anti-inflammatory agents. U.S. Patent Nos. 6,114,367 and 5,869,511 (National stage of WO 95/24398) disclose isoxazoline compounds as inhibitors of TNF release. WO 95/14681 discloses a series of isoxazoline compounds as anti-inflammatory agents. WO 02/100332 discloses isoxazoline compounds having macrophage inhibitory factor (MIF) antagonist activity.

Summary of the Invention

The present invention provides isoxazoline derivatives and their analogues, which can be used for the treatment of AIDS, asthma, arthritis, bronchitis, chronic obstructive pulmonary disease (COPD), psoriasis, allergic rhinitis, shock, atopic dermatitis, Crohn's disease, adult respiratory distress syndrome (ARDS), eosinophilic granuloma, allergic conjunctivitis, osteoarthritis, ulcerative colitis and other inflammatory diseases, and the processes for the synthesis of these compounds.

Pharmaceutically acceptable salts, pharmaceutically acceptable solvates, enantiomers, diastereomers or N-oxides of these compounds having the same type of activity are also provided.

Pharmaceutical compositions containing the compounds, which may also contain pharmaceutically acceptable carriers or diluents, can be used for the treatment of AIDS, asthma, arthritis, bronchitis, chronic obstructive pulmonary disease (COPD), psoriasis, allergic rhinitis, shock, atopic dermatitis, Crohn's disease, adult respiratory distress syndrome, eosinophilic granuloma, allergic conjunctivitis, osteoarthritis, ulcerative colitis and other inflammatory diseases.


WO 2005/051931

PCT/IB2004/003893

3

Other aspects will be set forth in the accompanying description which follows and in part will be apparent from the description or may be learnt by the practice of the invention.

In accordance with one aspect, there are provided compounds having the structure of Formula I:



Formula I

their pharmaceutically acceptable salts, pharmaceutically acceptable solvates, enantiomers, diastereomers or N-oxides.

When X is oxygen in Formula I:

- 15 R₁ can represent: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl; cyano; nitro; amino; substituted amino; hydroxyl; alkoxy; aryloxy; COR' or COOR' (wherein R' can be hydrogen, alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heterocyclyl, (heterocyclyl)alkyl, or (heteroaryl)alkyl); aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl) alkyl; (heterocyclyl) alkyl; (CH₂)₁₋₄OR' (wherein R' is as defined above, but
- 20 also including hydroxy); C(=O)NR_xR_y (wherein R_x and R_y can be independently selected from hydrogen, alkyl, C₃₋₆ alkenyl, C₃₋₆ alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heteroaryl, heterocyclyl, heteroarylalkyl, or heterocyclylalkyl); or (CH₂)_m-C(=O)R₃ [wherein m is an integer in the range of 0-2 and R₃ can be optionally substituted R_p or R_q (wherein R_p can be a 4-12 membered (un)saturated monocyclic or bicyclic ring containing
- 25 1-4 heteroatom(s) selected from N, O and S wherein the ring can be attached to (CH₂)_mC(=O) through N and R_q can be a 4-12 membered (un)saturated monocyclic or bicyclic ring containing 0-4 heteroatom(s) selected from the group consisting of N, O and S wherein the ring can be attached to (CH₂)_mC(=O) through C) and wherein the substituents of R₃ can be one or more of: alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl,
- 30 halogen, hydroxyl, alkoxy, aryloxy, nitro, cyano, amino, substituted amino, hydroxyalkyl,

WO 2005/051931

PCT/IB2004/003893

4

- oxo, acyl, optionally substituted amino (wherein the substituents are selected from C₁-C₆ alkyl, aryl, aralkyl, or cycloalkyl), aryl, carboxyl, alkaryl, carbamoyl, alkyl ether, C(=O)NR₅R₆ (wherein R₅ and R₆ are independently selected from hydrogen, alkyl, C₃₋₆ alkenyl, C₃₋₆ alkynyl, aryl, and aralkyl), optionally substituted monocyclic or bicyclic 4-12
- 5 membered carbocyclic ring system (wherein the optional substituent(s) is/are selected from alkyl, alkenyl, alkynyl, halogen, hydroxyl, and alkoxy), heteroaryl, heterocyclyl, heteroarylalkyl, or heterocyclalkyl].

- R₂ can represent: cyano; heteroaryl; heterocyclyl; or (CH₂)_nNHCOR₇ (wherein n represents an integer 1 to 6 and R₇ can represent hydrogen, alkyl, alkenyl, alkynyl,
- 10 (un)saturated, cycloalkyl, alkoxy, aryloxy, aryl, aralkyl, heteroaryl, heterocyclyl, (CH₂)₁₋₄OR' wherein R' is the same as defined above, or NR_xR_y wherein R_x and R_y are the same as defined above).

R₄ can represent: hydrogen; alkyl; halogen; cyano; carboxy; or C(=O)NR_xR_y wherein R_x and R_y are the same as defined above.

- 15 X₁ and X₂ can be independently selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl; acyl; aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl)alkyl; or (heterocyclyl)alkyl.

- Y can represent: an oxygen atom; a sulphur atom; or NR (wherein R is selected from hydrogen, alkyl, alkenyl, alkynyl, un(saturated) cycloalkyl, acyl, aryl, aralkyl, heteroaryl,
- 20 heterocyclyl, (heteroaryl)alkyl, or (heterocyclyl)alkyl).

Y₁ and Y₂ can be independently selected from: hydrogen; alkyl; nitro; cyano; halogen; OR wherein R is the same as defined earlier; SR wherein R is the same as defined earlier; NHR wherein R is the same as defined earlier; COOR'; or COR' wherein R' is the same as defined above.

- 25 Further, Y₁ and X₂, X₁ and Y₂, X₁ and X₂ may together form a cyclic ring fused with the ring A containing 3-5 carbon atoms within the ring and having 1-3 heteroatoms selected from N, O or S.

When X is NR₈ or S wherein R₈ is hydrogen, lower alkyl (C₁-C₆) or aryl:

R₁, R₄, X₁, X₂, Y, Y₁ and Y₂ are the same as defined above.


WO 2005/051931

PCT/TB2004/003893

5

R_2 can represent: $(CH)_nNHCOR_7$ (wherein n represents an integer 1 to 6 and R_7 is the same as defined above), with the proviso that when R_2 is heterocyclyl, R_1 can not be $(CH_2)_1-4OR'$, $C(=O)NR_xR_y$ or $(CH_2)_m-C(=O)R_3$.

In one particular embodiment, there are provided compounds having the structure
5 of Formula XXXII,



their pharmaceutically acceptable salts, pharmaceutically acceptable solvates, enantiomers, diastereomers or N-oxides.

15 In such compounds of Formula XXXII,

Y , Y_1 , Y_2 , R_1 and R_4 can be as defined for Formula I;

X_1 can be alkyl;

X_2 can be alkyl, cycloalkyl, or aralkyl;

X_3 , X_4 , X_5 and X_6 can be independently selected from C, CH, CH_2 , CO, CS, NH, N, O
20 and S;

R_{15} , R_{16} , and R_{17} can be independently selected from no atom, alkyl, $COCH_3$, $COOC_2H_5$, NH_2 , NH-cyclopropyl, CN and SH; and


----- represents an optional double bond.

25 In another embodiment, there are provided compounds having the structure of Formula XXXIII,

WO 2005/051931

PCT/IB2004/003893

6



their pharmaceutically acceptable salts, pharmaceutically acceptable solvates,

10 enantiomers, diastereomers or N-oxides.


In such compounds of Formula XXXIII,

Y, Y₁, Y₂, X₁, X₂, R₁ and R₄ can be as defined for Formula I;

X₇ can be O or S; and

R₁₈ can represent hydrogen, alkyl, aryl, heteroaryl, cycloalkyl, or heterocyclyl.

15 In yet another embodiment, there are provided compounds having the structure of Formula XXXIV,



their pharmaceutically acceptable salts, pharmaceutically acceptable solvates,
enantiomers, diastereomers or N-oxides.

25 In such compounds of Formula XXXIV,

Y, Y₁, Y₂, X₁, X₂, R₁ and R₄ can be as defined for Formula I; and

R₁₉ can represent cyano, -CONHNH₂, -C(NH₂)=N-O-C(O)R', (CH₂)_nNHCOR₇ or 6-membered heteroaryl, wherein R', R₇ and n are the same as defined for Formula I.

The following definitions apply to terms as used herein:

WO 2005/051931

PCT/IB2004/003893

7

The term "alkyl," unless otherwise specified, refers to a monoradical branched or unbranched saturated hydrocarbon chain having from 1 to 20 carbon atoms. This term can be exemplified by groups such as methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, t-butyl, n-pentyl, isopentyl, neopentyl, n-hexyl, n-decyl, tetradecyl, and the like.

- 5 Alkyl groups may be substituted further with one or more substituents selected from alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, oxo, thiocarbonyl, carboxy, carboxyalkyl, aryl, heterocyclyl, heteroaryl, arylthio, thiol, alkylthio, aryloxy, nitro, aminosulfonyl, aminocarbonylamino, -NHC(=O)R_f , $\text{-NR}_f\text{R}_q$, $\text{-C(=O)NR}_f\text{R}_q$,
10 $\text{-NHC(=O)NR}_f\text{R}_q$, -C(=O)heteroaryl , C(=O)heterocyclyl , $\text{-O-C(=O)NR}_f\text{R}_q$ {wherein R_f and R_q are independently selected from alkyl, alkenyl, cycloalkyl, cycloalkenyl, aryl, aralkyl, heterocyclyl, heteroaryl, heterocyclylalkyl, heteroarylalkyl}, nitro, or $\text{-SO}_2\text{R}_6$ (wherein R_6 is alkyl, alkenyl, alkynyl, cycloalkyl, aralkyl, aryl, heterocyclyl, heteroaryl, heteroarylalkyl or heterocyclylalkyl). Unless otherwise constrained by the definition,
15 alkyl substituents may be further substituted by 1-3 substituents selected from alkyl, carboxy, $\text{-NR}_f\text{R}_q$, $\text{-C(=O)NR}_f\text{R}_q$, $\text{-OC(=O)NR}_f\text{R}_q$, $\text{-NHC(=O)NR}_f\text{R}_q$ (wherein R_f and R_q are the same as defined earlier), hydroxy, alkoxy, halogen, CF_3 , cyano, and $\text{-SO}_2\text{R}_6$, (wherein R_6 are the same as defined earlier); or an alkyl group also may be interrupted by 1-5 atoms of groups independently selected from oxygen, sulfur or -NR_a - {wherein R_a is
20 selected from hydrogen, alkyl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, aryl, acyl, aralkyl, -C(=O)OR_f (wherein R_f is the same as defined earlier), SO_2R_6 (where R_6 is as defined earlier), or $\text{-C(=O)NR}_f\text{R}_q$ (wherein R_f and R_q are as defined earlier)}. Unless otherwise constrained by the definition, all substituents may be substituted further by 1-3 substituents selected from alkyl, carboxy, $\text{-NR}_f\text{R}_q$, $\text{-C(=O)NR}_f\text{R}_q$, $\text{-O-C(=O)NR}_f\text{R}_q$
25 (wherein R_f and R_q are the same as defined earlier) hydroxy, alkoxy, halogen, CF_3 , cyano, and $\text{-SO}_2\text{R}_6$ (where R_6 is same as defined earlier); or an alkyl group as defined above that has both substituents as defined above and is also interrupted by 1-5 atoms or groups as defined above.

- 30 The term "alkenyl," unless otherwise specified, refers to a monoradical of a branched or unbranched unsaturated hydrocarbon group having from 2 to 20 carbon atoms with cis, trans, or geminal geometry. In the event that alkenyl is attached to a heteroatom, the double bond cannot be alpha to the heteroatom. Alkenyl groups may be substituted

WO 2005/051931

PCT/IB2004/003893

8

further with one or more substituents selected from alkyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, -NHC(=O)R_f, -NR_fR_q, -C(=O)NR_fR_q, -NHC(=O)NR_fR_q, -O-C(=O)NR_fR_q (wherein R_f and R_q are the same as defined earlier), alkoxycarbonylamino, azido, cyano, halogen, hydroxy, oxo, thiocarbonyl, carboxy, arylthio, thiol, alkylthio, aryl, aralkyl, aryloxy, heterocyclyl, heteroaryl, heterocyclyl alkyl, heteroaryl alkyl, aminosulfonyl, aminocarbonylamino, alkoxyamino, nitro, or SO₂R₆ (wherein R₆ are is same as defined earlier). Unless otherwise constrained by the definition, alkenyl substituents optionally may be substituted further by 1-3 substituents selected from alkyl, carboxy, hydroxy, alkoxy, halogen, -CF₃, cyano, -NR_fR_q, -C(=O)NR_fR_q, -O-C(=O)NR_fR_q (wherein R_f and R_q are the same as defined earlier) and -SO₂R₆ (where R₆ is same as defined earlier).

The term "alkynyl," unless otherwise specified, refers to a monoradical of an unsaturated hydrocarbon, having from 2 to 20 carbon atoms. In the event that alkynyl is attached to a heteroatom, the triple bond cannot be alpha to the heteroatom. Alkynyl groups may be substituted further with one or more substituents selected from alkyl, alkenyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, oxo, thiocarbonyl, carboxy, arylthio, thiol, alkylthio, aryl, aralkyl, aryloxy, aminosulfonyl, aminocarbonylamino, nitro, heterocyclyl, heteroaryl, heterocyclylalkyl, heteroarylalkyl, -NHC(=O)R_f, -NR_fR_q, -NHC(=O)NR_fR_q, -C(=O)NR_fR_q, -O-C(=O)NR_fR_q (wherein R_f and R_q are the same as defined earlier), or -SO₂R₆ (wherein R₆ is as defined earlier). Unless otherwise constrained by the definition, alkynyl substituents optionally may be substituted further by 1-3 substituents selected from alkyl, carboxy, carboxyalkyl, hydroxy, alkoxy, halogen, CF₃, -NR_fR_q, -C(=O)NR_fR_q, -NHC(=O)NR_fR_q, -C(=O)NR_fR_q (wherein R_f and R_q are the same as defined earlier), cyano, or -SO₂R₆ (where R₆ is same as defined earlier).

The term "cycloalkyl," unless otherwise specified, refers to cyclic alkyl groups of from 3 to 20 carbon atoms having a single cyclic ring or multiple condensed rings, which may optionally contain one or more olefinic bonds, unless otherwise constrained by the definition. Such cycloalkyl groups can include, for example, single ring structures, including cyclopropyl, cyclobutyl, cyclooctyl, cyclopentenyl, and the like, or multiple ring structures, including adamantanyl, and bicyclo [2.2.1] heptane, or cyclic alkyl groups to which is fused an aryl group, for example, indane, and the like. Spiro and fused ring

WO 2005/051931

PCT/IB2004/003893

9

structures can also be included. Cycloalkyl groups may be substituted further with one or more substituents selected from alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, oxo, thiocarbonyl, carboxy, carboxyalkyl, arylthio, thiol, alkylthio, aryl, aralkyl, aryloxy, aminosulfonyl, aminocarbonylamino, $-NR_fR_q$, $-NHC(=O)NR_fR_q$, $-NHC(=O)R_f$, $-C(=O)NR_fR_q$, $-O-C(=O)NR_fR_q$ (wherein R_f and R_q are the same as defined earlier), nitro, heterocyclyl, heteroaryl, heterocyclylalkyl, heteroarylalkyl, or SO_2-R_6 (wherein R_6 is same as defined earlier). Unless otherwise constrained by the definition, cycloalkyl substituents optionally may be substituted further by 1-3 substituents selected from alkyl, carboxy, hydroxy, alkoxy, halogen, CF_3 , $-NR_fR_q$, $-C(=O)NR_fR_q$, $-NHC(=O)NR_fR_q$, $-O-C(=O)NR_fR_q$ (wherein R_f and R_q are the same as defined earlier), cyano or $-SO_2R_6$ (where R_6 is same as defined earlier).

The term "alkoxy" denotes the group O-alkyl, wherein alkyl is the same as defined above.

The term "aryl," unless otherwise specified, refers to carbocyclic aromatic groups, for example, phenyl, biphenyl or naphthyl ring and the like, optionally substituted with 1 to 3 substituents selected from halogen (e.g., F, Cl, Br, I), hydroxy, alkyl, alkenyl, alkynyl, cycloalkyl, alkoxy, acyl, aryloxy, CF_3 , cyano, nitro, $COOR_e$ (wherein R_e is hydrogen, alkyl, alkenyl, cycloalkyl, aralkyl, heterocyclylalkyl, heteroarylalkyl), $NHC(=O)R_f$, $-NR_fR_q$, $-C(=O)NR_fR_q$, $-NHC(=O)NR_fR_q$, $-O-C(=O)NR_fR_q$ (wherein R_f and R_q are the same as defined earlier), $-SO_2R_6$ (wherein R_6 is same as defined earlier), carboxy, heterocyclyl, heteroaryl, heterocyclylalkyl, heteroarylalkyl or amino carbonyl amino. The aryl group optionally may be fused with a cycloalkyl group, wherein the cycloalkyl group may optionally contain heteroatoms selected from O, N or S.

The term "aralkyl," unless otherwise specified, refers to alkyl-aryl linked through an alkyl portion (wherein alkyl is as defined above) and the alkyl portion contains 1-6 carbon atoms and aryl is as defined below. Examples of aralkyl groups include benzyl, ethylphenyl and the like.

The term "aralkenyl," unless otherwise specified, refers to alkenyl-aryl linked through alkenyl (wherein alkenyl is as defined above) portion and the alkenyl portion contains 1 to 6 carbon atoms and aryl is as defined below.

The term "aryloxy" denotes the group O-aryl, wherein aryl is as defined above.

WO 2005/051931

PCT/IB2004/003893

10

The term "carboxy," as defined herein, refers to $-C(=O)OH$.

The term "heteroaryl," unless otherwise specified, refers to an aromatic ring structure containing 5 or 6 ring atoms, or a bicyclic aromatic group having from 8 to 10 ring atoms, with one or more heteroatom(s) independently selected from N, O or S optionally substituted with 1 to 4 substituent(s) selected from halogen (*e.g.*, F, Cl, Br, I), hydroxy, alkyl, alkenyl, alkynyl, cycloalkyl, acyl, carboxy, aryl, alkoxy, aralkyl, cyano, nitro, heterocyclyl, heteroaryl, $-NR_fR_q$, $CH=NOH$, $-(CH_2)_wC(=O)R_g$ {wherein *w* is an integer from 0-4 and R_g is hydrogen, hydroxy, OR_f , NR_fR_q , $-NHOR_z$ or $-NHOH$ }, $-C(=O)NR_fR_q$ and $-NHC(=O)NR_fR_q$, $-SO_2R_6$, $-O-C(=O)NR_fR_q$, $-O-C(=O)R_f$, $-O-C(=O)OR_f$ (wherein R_6 , R_f and R_q are as defined earlier, and R_z is alkyl, cycloalkyl, aryl, heteroaryl, heterocyclyl, heteroarylalkyl or heterocyclylalkyl). Unless otherwise constrained by the definition, the substituents are attached to a ring atom, *i.e.*, carbon or heteroatom in the ring. Examples of heteroaryl groups include oxazolyl, imidazolyl, pyrrolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, tetrazolyl, thiazolyl, oxadiazolyl, benzoimidazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, thienyl, isoxazolyl, triazinyl, furanyl, benzofuranyl, indolyl, benzothiazolyl, or benzoxazolyl, and the like.

The term "heterocyclyl," unless otherwise specified, refers to a non-aromatic monocyclic or bicyclic cycloalkyl group having 5 to 10 atoms wherein 1 to 4 carbon atoms in a ring are replaced by heteroatoms selected from O, S or N, and optionally are benzofused or fused heteroaryl having 5-6 ring members and/or optionally are substituted, wherein the substituents are selected from halogen (*e.g.*, F, Cl, Br, I), hydroxy, alkyl, alkenyl, alkynyl, cycloalkyl, acyl, aryl, alkoxy, alkaryl, cyano, nitro, oxo, carboxy, heterocyclyl, heteroaryl, $-O-C(=O)R_f$, $-O-C(=O)OR_f$, $-C(=O)NR_fR_q$, SO_2R_6 , $-O-C(=O)NR_fR_q$, $-NHC(=O)NR_fR_q$, $-NR_fR_q$ (wherein R_6 , R_f and R_q are as defined earlier) or guanidine. Heterocyclyl can optionally include rings having one or more double bonds. Unless otherwise constrained by the definition, the substituents are attached to the ring atom, *i.e.*, carbon or heteroatom in the ring. Also, unless otherwise constrained by the definition, the heterocyclyl ring optionally may contain one or more olefinic bond(s). Examples of heterocyclyl groups include oxazolidinyl, tetrahydrofuranyl, dihydrofuranyl, dihydropyridinyl, dihydroisoxazolyl, dihydrobenzofuryl, azabicyclohexyl, dihydroindolyl, pyridinyl, isoindole 1,3-dione, piperidinyl or piperazinyl.

WO 2005/051931

PCT/IB2004/003893

11

"Heteroarylalkyl" refers to alkyl-heteroaryl group linked through alkyl portion, wherein the alkyl and heteroaryl are as defined earlier.

"Heterocyclalkyl" refers to alkyl-heterocyclalkyl group linked through alkyl portion, wherein the alkyl and heterocyclalkyl are as defined earlier.

5 "Acyl" refers to $-C(=O)R''$ wherein R'' is selected from hydrogen, alkyl, cycloalkyl, aryl, aralkyl, heteroaryl, heterocyclalkyl, heteroarylalkyl or heterocyclalkyl.

"Alkylcarbonyl" refers to $-C(=O)R''$, wherein R'' is selected from alkyl, cycloalkyl, aryl, aralkyl, heteroaryl, heterocyclalkyl, heteroarylalkyl or heterocyclalkyl.

10 "Alkylcarboxy" refers to $-O-C(=O)R''$, wherein R'' is selected from alkyl, cycloalkyl, aryl, aralkyl, heteroaryl, heterocyclalkyl, heteroarylalkyl or heterocyclalkyl.

"Amine," unless otherwise specified, refers to $-NH_2$. "Substituted amine," unless otherwise specified, refers to $-N(R_k)_2$, wherein each R_k independently is selected from hydrogen {provided that both R_k groups are not hydrogen (defined as "amino")}, alkyl, alkenyl, alkynyl, aralkyl, cycloalkyl, aryl, heteroaryl, heterocyclalkyl, heterocyclalkyl, 15 heteroarylalkyl, acyl, SO_2R_6 (wherein R_6 is as defined above), $-C(=O)NR_fR_q$, $NHC(=O)NR_fR_q$, or $-NHC(=O)OR_f$ (wherein R_f and R_q are as defined earlier),

"Thiocarbonyl" refers to $-C(=S)H$. "Substituted thiocarbonyl" refers to $-C(=S)R''$, wherein R'' is selected from alkyl, cycloalkyl, aryl, aralkyl, heteroaryl, heterocyclalkyl, heteroarylalkyl or heterocyclalkyl, amine or substituted amine.

20 Unless otherwise constrained by the definition, all substituents optionally may be substituted further by 1-3 substituents selected from alkyl, aralkyl, cycloalkyl, aryl, heteroaryl, heterocyclalkyl, carboxy, carboxyalkyl, hydroxy, alkoxy, halogen, CF_3 , cyano, $-C(=T)NR_fR_q$, $-O(C=O)NR_fR_q$ (wherein R_f , R_q and T are the same as defined earlier) and $-OC(=T)NR_fR_q$, $-SO_2R_6$ (where R_6 is the same as defined earlier).

25 The term "leaving group" refers to groups that exhibit or potentially exhibit the properties of being labile under the synthetic conditions and also, of being readily separated from synthetic products under defined conditions. Examples of leaving groups include, but are not limited to, halogen (e.g., F, Cl, Br, I), triflates, tosylate, mesylates, alkoxy, thioalkoxy, or hydroxy radicals and the like.

30 The term "protecting groups" refers to moieties that prevent chemical reaction at a location of a molecule intended to be left unaffected during chemical modification of such

WO 2005/051931

PCT/IB2004/003893

12

molecule. Unless otherwise specified, protecting groups may be used on groups, such as hydroxy, amino, or carboxy. Examples of protecting groups are found in T.W. Greene and P.G.M. Wuts, "Protective Groups in Organic Synthesis", 2nd Ed., John Wiley and Sons, New York, N.Y., which is incorporated herein by reference. The species of the carboxylic protecting groups, amino protecting groups or hydroxy protecting groups employed are not critical, as long as the derivatised moieties/moiety is/are stable to conditions of subsequent reactions and can be removed without disrupting the remainder of the molecule.

The term "pharmaceutically acceptable salts" refers to derivatives of compounds that can be modified by forming their corresponding acid or base salts. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acids salts of basic residues (such as amines), or alkali or organic salts of acidic residues (such as carboxylic acids), and the like.

The compounds provided herein can be used for treating AIDS, asthma, arthritis, bronchitis, chronic obstructive pulmonary disease, psoriasis, allergic rhinitis, shock, atopic dermatitis, Crohn's disease, adult respiratory distress syndrome, eosinophilic granuloma, allergic conjunctivitis, osteoarthritis, ulcerative colitis and other inflammatory diseases.

In accordance with yet another aspect, there are provided processes for the preparation of the compounds as described herein.

20


Detailed Description of the Invention

The compounds described herein may be prepared by techniques well known in the art and familiar to the average synthetic organic chemist. In addition, the compounds of present invention may be prepared by the following reaction sequences as depicted in schemes I, IA, IB, II, III, IV and V.

WO 2005/051931

PCT/IB2004/003893

13



10 The compounds of Formula VII (a) can be prepared according to Scheme I. Thus, reacting a compound of Formula II with compound of Formula X_2Z (wherein Z is halogen) to give a compound of Formula III [wherein X_1 , X_2 (except hydrogen), Y_1 and Y_2 are the same as defined earlier], which on reaction with hydroxylamine hydrochloride gives a compound of Formula IV, which on treatment with a compound of Formula V

15 gives a compound of Formula VI (wherein R_1 and R_4 are the same as defined earlier and R_r represents $[(CH_2)_nCN, COOH, COOCH_3, CHO$ or pyridyl, wherein n is 0 to 2)], which on reaction with hydroxylamine hydrochloride (when R_r is CN) to give a compound of Formula VII, which is finally reacted with a compound of Formula $(R'CO)_2O$ to give a compound of Formula VII(a) (wherein R' is the san as defined earlier) .

20 The reaction of a compound of Formula II with a compound of Formula X_2Z to give a compound of Formula III can be carried out in a solvent, for example, tetrahydrofuran, dimethylformamide, dimethylsulphoxide or acetonitrile.

25 The reaction of a compound of Formula II with compound of Formula X_2Z can be carried out in the presence of potassium iodide and an inorganic base, for example, sodium carbonate, sodium bicarbonate, potassium carbonate or potassium bicarbonate.

 The reaction of a compound of Formula III with hydroxylamine hydrochloride to give a compound of Formula IV can be carried out in the presence of sodium acetate or potassium acetate in a solvent , for example, methanol, ethanol, propanol or n-butanol.

30 The reaction of a compound of Formula IV with a compound of Formula V to give a compound of Formula VI can be carried out in the presence of sodium hypochlorite in a

WO 2005/051931

PCT/IB2004/003893

14

solvent, for example, tetrahydrofuran, dimethylformamide, dimethylsulphoxide or acetonitrile.

The reaction of a compound of Formula VI with hydroxylamine hydrochloride to give a compound of Formula VII can be carried out in a solvent, for example,


- 5 tetrahydrofuran, dimethylformamide, dimethylsulphoxide, acetonitrile, acetone, ethanol or mixtures thereof.

The reaction of a compound of Formula VI with hydroxylamine hydrochloride can be carried out in the presence of an inorganic base, for example, sodium carbonate, sodium bicarbonate, potassium carbonate or potassium bicarbonate.

- 10 The reaction of a compound of Formula VII with a compound of Formula $(R'CO)_2O$ to give a compound of Formula VII (a) can be carried out in a solvent, for example, tetrahydrofuran, dimethylformamide, dimethylsulphoxide or acetonitrile.

The reaction of a compound of Formula VII with a compound of Formula $(R'CO)_2O$ can be carried out in the presence of an organic base, for example,

- 15 trimethylamine, triethylamine or pyridine.



The compounds of Formula IX and X can be prepared according to scheme 1A.

- 25 Thus, reacting a compound of

- (a) Formula VI (when R_r is $COOCH_3$) with hydrazine hydrate to give a compound of Formula VIII (wherein X_1 , X_2 , Y_1 , Y_2 , R_1 and R_4 are the same as defined earlier), which on reaction with a compound of Formula $HC(OR_{11})_3$ gives a compound of Formula IX (wherein R_{11} represents alkyl from C_1 to C_3); or

WO 2005/051931

PCT/IB2004/003893

15

(b) Formula VI (when Rr is CN) with sodium azide to give a compound of Formula X (wherein X₁, X₂, Y₁, Y₂, R₁ and R₄ are the same as defined earlier),

The reaction of a compound of Formula VI with hydrazine hydrate to give a compound of Formula VIII can be carried out at a temperature ranging, for example, from 120 to 140°C.

The reaction of a compound of Formula VIII with a compound of Formula HC(OR₁₁)₃ to give a compound of Formula IX can be carried out at a temperature ranging, for example, from 120 to 160°C.

The reaction of a compound of Formula VI with sodium azide to give a compound of Formula X can be carried out in a solvent, for example, benzene, toluene or xylene.


The reaction of a compound of Formula VI with sodium azide to give a compound of Formula X can be carried out in the presence of hydrochloride salt of an organic base, for example, trimethylamine, triethylamine or pyridine.

Scheme IB

15

20

25



WO 2005/051931

PCT/IB2004/003893

16

The compounds of Formulae XI-XV can be prepared according to scheme IB.

Thus, reacting a compound of Formula VII (wherein X_1 , X_2 , Y_1 , Y_2 , R_1 and R_4 are the same as defined earlier) with

- (a) methyl chloroformate to give a compound of Formula XI;
- 5 (b) thiocarbonyl diimidazole and 1,8-diazabicyclo[5.4.0]undec-7-one to give a compound of Formula XII, which on treatment with a compound of Formula $R_{11}Z$ (wherein Z is halogen) gives a compound of Formula XIII (wherein R_{11} is alkyl);
- (c) thiocarbonyl diimidazole and boron trifluoride etherate to give a compound of Formula XIV;
- 10 (d) a compound of Formula $R_{12}COOH$,
- (e) a compound of Formula $R_{12}COCl$ or
- (f) a compound of Formula $R_{12}COOC_2H_5$ to give a compound of Formula XV (wherein R_{12} is alkyl, cycloalkyl, aryl, heteroaryl or heterocyclyl).

15 The reaction of a compound of Formula VII with methyl chloroformate to give a compound of Formula XI can be carried out in a solvent, for example, xylene, benzene or toluene.

The reaction of a compound of Formula VII with methyl chloroformate can be carried out in the presence of an organic base, for example, pyridine, trimethylamine or triethylamine.

20 The reaction of a compound of Formula VII with thiocarbonyl diimidazole and 1,8-diazabicyclo [5.4.0] undec-7-one to give a compound of Formula XII can be carried out in a solvent, for example, acetonitrile, acetone, dimethylformamide, dimethylsulfoxide or tetrahydrofuran.

25 The reaction of a compound of Formula XII with a compound of Formula $R_{11}Z$ to give a compound of Formula XIII can be carried out in a solvent, for example, acetone, acetonitrile, tetrahydrofuran or dimethylformamide.

The reaction of a compound of Formula XII with a compound of Formula $R_{11}Z$ can be carried out in the presence of an inorganic base, for example, sodium carbonate, sodium bicarbonate, potassium carbonate or potassium bicarbonate.

WO 2005/051931

PCT/IB2004/003893

17


The reaction of a compound of Formula VII with a compound of Formula $R_{12}COOH$ to give a compound of Formula XV can be carried out in the presence of isobutylchloroformate and an organic base, for example, triethylamine, dimethylamine or pyridine in a solvent, for example, dimethylformamide, tetrahydrofuran or acetonitrile.

- 5 The reaction of a compound of Formula VII to give a compound of Formula XV can be carried out in the presence of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride, 1-hydroxybenzotriazole and N-methylmorpholine.

- The reaction of a compound of Formula VII with a compound of Formula $R_{12}COCl$ to give a compound of Formula XV can be carried out in a solvent, for example, 10 toluene, acetonitrile, acetone, dimethylformamide, dimethylsulphoxide or tetrahydrofuran.

The reaction of a compound of Formula VII with a compound of Formula $R_{12}COOC_2H_5$ to give a compound of Formula XV can be carried out in the presence of an inorganic base, for example, sodium carbonate, potassium carbonate or sodium hydride in a solvent, for example, dimethylformamide, tetrahydrofuran or acetonitrile.

- 15 The reaction of a compound of Formula VII with the compounds of Formula $R_{12}COOH$, $R_{12}COCl$ and $R_{12}COOC_2H_5$ can be carried out in the presence of molecular sieves.




- The compounds of Formula XVb can be prepared according to Scheme IC. Thus reacting a compound of Formula XVa with 2-oxo propionic acid ethyl ester gives a 25 compound of Formula XVb (wherein X_1 , X_2 , Y_1 , Y_2 , R_1 , and R_4 are the same as earlier). The reaction can be carried out in a solvent, for example, acetonitrile, acetone, dimethylformamide, dimethylsulphoxide or tetrahydrofuran.

WO 2005/051931

PCT/IB2004/003893

18



The compounds of Formula XX can be prepared according to Scheme II. Thus reacting a compound of Formula IV with a compound of Formula XVI to give a compound of Formula XVII (wherein X_1 , X_2 , Y_1 , Y_2 , R_1 , R_4 , Z and n are the same as defined earlier), which on treatment with potassium phthalamide gives a compound of Formula XVIII, which on treatment with a hydrazine hydrate gives a compound of Formula XIX, which is finally treated with a compound of Formula $R_{12}COCl$ or $R_{12}COOH$ to give a compound of Formula XX (wherein R_{12} is the same as defined earlier).

The reaction of a compound of Formula IV with a compound of Formula XVI to give a compound of Formula XVII can be carried out in a solvent, for example, acetonitrile, acetone, dimethylformamide, dimethylsulphoxide or tetrahydrofuran.

The reaction of a compound of Formula XVII with potassium phthalamide to give a compound of Formula XVIII can be carried out in a solvent, for example, acetonitrile, acetone, dimethylformamide, dimethylsulphoxide or tetrahydrofuran.

The reaction of a compound of Formula XVIII with hydrazine hydrate to give a compound of Formula XIX can be carried out in a solvent, for example, methanol, ethanol, propanol, butanol, water or mixture thereof.

The reaction of a compound of Formula XIX with a compound of Formula $R_{12}COCl$ to give a compound of Formula XX can be carried out in a solvent, for example, chloroform, dichloromethane or dichloroethane.


WO 2005/051931

PCT/IB2004/003893

19

The reaction of a compound of Formula XIX with a compound of Formula $R_{12}COCl$ can be carried out in the presence of an organic base, for example, trimethylamine, triethylamine or pyridine.

The reaction of a compound of Formula XIX with a compound of Formula $R_{12}COOH$ to give a compound of Formula XX can be carried out in the presence of 1-ethyl-3-[3-(dimethylamino)propyl]carbodiimide, 1-hydroxybenzotriazole and N-methylmorpholine in a solvent, for example, dimethylformamide, dimethylsulfoxide or tetrahydrofuran.



The compounds of Formula XXIII can be prepared according to Scheme III. Thus, reacting a compound of Formula XXI with hydroxylamine hydrochloride to give a compound of Formula XXII (wherein R_{13} is alkyl, aryl or heteroaryl), which on reaction with a compound of Formula VI (when R_r is $COOH$, scheme I) gives a compound of Formula XXIII (wherein X_1 , X_2 , Y_1 , Y_2 , R_1 and R_4 are the same as defined earlier).


The reaction of a compound of Formula XXI to give a compound of Formula XXII can be carried out in the presence of sodium carbonate or potassium carbonate in a solvent, for example, methanol, ethanol, propanol, n-butanol, water or mixture thereof.

The reaction of a compound of Formula XXII with a compound of Formula VI to give a compound of Formula XXIII can be carried out in a solvent, for example, dimethylformamide or dimethylsulfoxide.

The reaction of a compound of Formula XXII with a compound of Formula VI can be carried out in the presence of 1-hydroxybenzotriazole, N-methylmorpholine and a coupling agent, for example, 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide hydrochloride or 1,3-dicyclohexyl carbodiimide.

The reaction of a compound of Formula XXII with a compound of Formula VI to give a compound of Formula XXIII can be carried out in the presence of sodium acetate or

potassium acetate solvent, for example, methanol, ethanol, propanol, n-butanol, water or mixture thereof.



The compounds of Formula XXIV-XXVII can be prepared according to scheme IV. Thus, reacting a compound of

- (1) Formula VI (when R_r is CN) with NH₂CH₂CH₂SH. HCl to give a compound of Formula XXIV (wherein X₁, X₂, Y₁, Y₂, R₁ and R₄ are the same as defined earlier);
- (2) Formula VI (when R_r is COOH) with NH₂NHCSNHR₁₄ to give a compound of Formula XXV (wherein X₁, X₂, Y₁, Y₂, R₁ and R₄ are the same as defined earlier, R₁₄ represents hydrogen, alkyl or cycloalkyl); or
- (3) Formula VI (when R_r is CHO) with hydroxylamine hydrochloride to give a compound of Formula XXVI which on reaction with methacrylonitrile gives a compound of Formula XXVII (wherein X₁, X₂, Y₁, Y₂, R₁ and R₄ are the same as defined earlier).

The reaction of a compound of Formula VI with NH₂CH₂CH₂SH. HCl to give a compound of Formula XXIV can be carried out in the presence of an organic base, for

WO 2005/051931

PCT/IB2004/003893


21

example, triethylamine, trimethylamine or pyridine in a solvent, for example, methanol, ethanol or isopropanol.

The reaction of a compound of Formula VI with $\text{NH}_2\text{NHCSNHR}_{14}$ to give a compound of Formula XXV can be carried out in the presence of POCl_3 in a solvent, for example, methanol or dioxane.

The reaction of a compound of Formula VI with hydroxylamine hydrochloride and sodium acetate to give a compound of Formula XXVI can be carried out in a solvent, for example, methanol, ethanol or isopropanol.

The reaction of a compound of Formula XXVI with methacrylonitrile to give a compound of Formula XXVII can be carried out in the presence of sodium hypochlorite in a solvent, for example, tetrahydrofuran, dimethylformamide, dimethylsulphoxide or acetonitrile.



The compounds of Formula XXIX-XXXI can be prepared according to Scheme V. Thus, reacting a compound of Formula VIII

- (1) with ethylmethylketone to give a compound of Formula XXVIII, which on treatment with acetic anhydride gives a compound of Formula XXIX (wherein X_1 , X_2 , Y_1 , Y_2 , R_1 and R_4 are the same as defined earlier)
- (2) with carbon disulphide to give a compound of Formula XXX, which on treatment with hydrazine hydrate gives a compound of Formula XXXI (wherein X_1 , X_2 , Y_1 , Y_2 , R_1 and R_4 are the same as defined earlier).

WO 2005/051931

PCT/IB2004/003893

22

The reaction of a compound of Formula VIII with ethylmethylketone to give a compound of Formula XXVIII can be carried out in a solvent, for example, methanol, ethanol or isopropanol.

5 The reaction of a compound of Formula XXVIII with acetic acid to give a compound of Formula XXIX can be carried out in the presence of an organic base, for example, pyridine, triethylamine or trimethylamine.

The reaction of a compound of Formula VIII with carbon disulphide to give a compound of Formula XXX can be carried out in the presence of an inorganic base, for example, sodium hydroxide, potassium hydroxide or calcium hydroxide.

10 The reaction of a compound of Formula VIII with carbon disulphide to give a compound of Formula XXX can be carried out in a solvent, for example, methanol, ethanol or isopropanol.

The reaction of a compound of Formula XXX with hydrazine hydrate to give a compound of Formula XXXI can be carried out in a solvent, for example, methanol,
15 ethanol or isopropanol.

In the above schemes, where the specific solvents, bases, coupling agents etc., are mentioned, it is to be understood that other solvents, bases coupling agents etc., known to those skilled in the art may be used. Similarly, the reaction temperature and duration may be adjusted according to the desired needs.

20 An illustrative list of compounds of the invention are listed below (also shown in Table 1 2, 3, 4, 5, 6 and 7)

-3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazol-5-yl]-4H-[1,2,4]oxadiazol-5-one (Compound No. 1),

25 -3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazol-5-yl]-4H-[1,2,4]oxadiazole-5-thione (Compound No. 2),

-3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazol-5-yl]-4H-[1,2,4]thiadiazol-5-one (Compound No. 3),

-2-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 4),

WO 2005/051931

PCT/IB2004/003893

23

- 2-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazol-5-yl]-5-methyl-[1,3,4]oxadiazole (Compound No. 5),
- 3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazol-5-yl]-4-methyl-4H-[1,2,4]oxadiazole-5-thione (Compound No. 6),
- 5 –3-{3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazol-5-yl]-[1,2,4]oxadiazol-5-yl}pyridine (Compound No. 7),
- 5-tert-Butyl-3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 8),
- 10 –5-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazol-5-yl]-1H-tetrazole (Compound No. 9),
- 3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazole-5-carbonitrile (Compound No. 10),
- Morpholine-4-carboxylic acid [3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazol-5-ylmethyl]amide (Compound No. 11),
- 15 –N-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazol-5-ylmethyl]-4-fluoro-benzamide (Compound No. 12),
- Adamantane-1-carboxylic acid [3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazol-5-yl-methyl]amide (Compound No. 13),
- 20 –Furan-2-carboxylic acid [3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazol-5-yl-methyl]amide (Compound No. 14),
- 2-(3-Cyclopentyloxy-4-methoxyphenyl)-N-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl]-acetamide (Compound No. 15),
- 1-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl-methyl]-3-(2-trifluoromethyl)-phenyl)-urea (Compound No. 16),
- 25 –1-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl-methyl]-3-(2,4-difluorophenyl)-urea (Compound No. 17),
- 1-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl-methyl]-3-O-tolyl-urea (Compound No. 18),

WO 2005/051931

PCT/IB2004/003893

24

- Morpholine-4-carboxylic acid [3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl-methyl]-amide (Compound No. 19),
- 3-(2-Chloro-6-trifluoromethylphenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl-[1,2,4]oxadiazole (Compound No. 20),
- 5 -3-(2-Chloro-4-fluorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 21),
- 3-(4-Chloro-2-methoxyphenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 22),
- 10 -3-(3-Chloro-4-fluorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 23),
- 3-(3-Chloro-4-methoxyphenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 24),
- 3-(3-Fluorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 25),
- 15 -3-(3,4-Difluorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazol-5-yl]- [1,2,4]oxadiazole (Compound No. 26),
- 3-(4-Methoxyphenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazol-5-yl]- [1,2,4]oxadiazole (Compound No. 27),
- 20 -3-(3,4-Dimethoxyphenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]- [1,2,4]oxadiazole (Compound No. 28),
- 3-(2-Chloro-6-fluorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 29),
- 3-(2,5-Difluorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]- [1,2,4]oxadiazole (Compound No. 30),
- 25 -3-(2,6-Dichlorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]- [1,2,4]oxadiazole (Compound No. 31),
- 3-(2,3-Dichlorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]- [1,2,4]oxadiazole (Compound No. 32),

WO 2005/051931

PCT/IB2004/003893

25

- 3-(2,4-Dichlorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 33),
- 3-(3,5-Dichlorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 34),
- 5 —3-(2,5-Dichlorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 35),
- 3-(3,5-Difluorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 36),
- 10 —3-(3-Chlorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 37),
- 3-(2,4-Difluoro-phenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 38),
- 3-(3,4-Dichlorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 39),
- 15 —3-(4-Chlorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 40),
- 4-{5-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole-3-yl}-phenylamine (Compound No. 41),
- 20 —3-Phenyl-5-[3-(3-cyclopentyloxy-4-methoxy-phenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 42),
- 3-(3,4-Dimethyl-phenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 43),
- 3-(2-Chlorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 44),
- 25 —3-(4-Fluorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 45),
- 3-Methyl-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 46),

WO 2005/051931

PCT/IB2004/003893

26

- 3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-5-oxazol-5-yl-4,5-dihydro-isoxazole (Compound No. 47),
- 5-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]thiadiazol-2-yl-amine (Compound No. 48),
- 5 -5-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]thiadiazol-2-yl}-cyclopropylamine (Compound No. 49),
- 1-{5-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-2-ethyl-2-methyl-[1,3,4]oxadiazol-3-yl}-ethanone (Compound No. 50),
- 10 -3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-(4,5-dihydro-thiazol-2-yl)-5-methyl-4,5-dihydro-isoxazole (Compound No. 51),
- 3'-(3-Cyclopentyloxy-4-methoxyphenyl)-5,5'-dimethyl-4,5,4',5'-tetrahydro-[3,5']biisoxazolyl-5-carbonitrile (Compound No. 52),
- 3-{3-[3-(cyclopentyloxy)-4-methoxyphenyl]-5-methyl-4,5-dihydroisoxazol-5-yl}-5-methyl-1,4,2-dioxazole-5-carboxylic acid ethyl ester (Compound No. 53)
- 15 -4-Amino-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-4H-[1,2,4]triazole-3-thiol (Compound No. 54),
- 5-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-3H-[1,3,4]oxadiazole-2-thione (Compound No. 55),
- 20 -3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-methyl-[1,2,4]oxadiazole (Compound No. 56),
- 4-{3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazol-5-yl}-pyridine (Compound No. 57),
- 5-tert-Butyl-3-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 58),
- 25 -3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-(2-ethoxy-phenyl)-[1,2,4]oxadiazole (Compound No. 59),
- 3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-cyclopropyl-[1,2,4]oxadiazole (Compound No. 60),

WO 2005/051931

PCT/IB2004/003893

27

-5-(3-Chlorophenyl)-3-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 61),

-5-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-m-tolyl-[1,2,4]oxadiazole (Compound No. 62),

5 -3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-(3,5-dimethyl-phenyl)-[1,2,4]oxadiazole (Compound No. 63),

-2,6-Dichloro-4-{3-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazol-5-yl}-pyridine (Compound No. 64),

10 -3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-isopropyl-[1,2,4]oxadiazole (Compound No. 65),

-5-Cyclohexyl-3-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 66),

-3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-fluoromethyl-[1,2,4]oxadiazole (Compound No. 67),

15 -3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-(tetrahydro-furan-2-yl)-[1,2,4]oxadiazole (Compound No. 68),

-3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-(4-fluoro-phenyl)-[1,2,4]oxadiazole (Compound No. 69),

20 -3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-(3-fluorophenyl) [1,2,4]oxadiazol (Compound No. 70),

-{3-[3-(3-Cyclopentyloxy-4-mthoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazol-5-yl}-acetonitrile (Compound No. 71),

-4-{3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole-5-yl}-benzonitrile (Compound No. 72),

25 -3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-trifluoromethyl-[1,2,4]oxadiazole (Compound No. 73),

-3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-(3-methoxy-phenyl)-[1,2,4]oxadiazole (Compound No. 74),

WO 2005/051931

PCT/IB2004/003893

28

- 3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-(3,4-dimethoxy-phenyl)-[1,2,4]oxadiazole (Compound No. 75),
- 5-(2-Chlorophenyl)-3-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 76),
- 5 -3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-(2,4-dichloro-phenyl)-[1,2,4]oxadiazole (Compound No. 77),
- 5-Cyclopentyl-3-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 78),
- 10 -3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-(3,4-dichloro-phenyl)-[1,2,4]oxadiazole (Compound No. 79),
- 2-[3-(4-Difluoromethoxy-3-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 80),
- 2-[3-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-difluoromethoxyphenyl]-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 81),
- 15 -2-[3-[3-benzyloxy-4-difluoromethoxyphenyl]-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 82),
- 2-[3-[4-Difluoromethoxy-3-ethoxyphenyl]-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 83),
- 2-[3-[4-Difluoromethoxy-3-isopropoxyphenyl]-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 84),
- 20 -2-[3-(3-Cyclohexyloxy-4-difluoromethoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 85),
- 2-[3-(3-Butoxy-4-difluoromethoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 86),
- 25 -2-[3-(3-Cycloheptyloxy-4-difluoromethoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 87),
- 2-[3-(4-Difluoromethoxy-3-propoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 88),

WO 2005/051931

PCT/IB2004/003893

29

- 2-[3-(3,4-Bis-difluoromethoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 89),
- 2-[3-(3-Cyclopentyloxy-4-difluoromethoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 90),
- 5 -2-[3-(3-Cyclopropylmethoxy-4-difluoromethoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 91),
- 2-[3-(3-Cyclopropylmethoxy-4-methoxy-phenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 92),
- 10 -2-[3-(3-Difluoromethoxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 93),
- 3-(3-Cyclopropylmethoxy-4-difluoromethoxyphenyl)-5-methyl-4,5-dihydro-isoxazole-5-carbonitrile (Compound No. 94),
- 3-(3-Cyclopentyloxy-4-difluoromethoxyphenyl)-5-methyl-4,5-dihydro-isoxazole-5-carbonitrile (Compound No. 95),
- 15 -3-(3,4-Bis-difluoromethoxyphenyl)-5-methyl-4,5-dihydro-isoxazole-5-carbonitrile (Compound No. 96),
- 3-(3-Cyclopropylmethoxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazole-5-carbonitrile (Compound No. 97),
- 20 -3-(3-Butoxy-4-difluoromethoxyphenyl)-5-methyl-4,5-dihydro-isoxazole-5-carbonitrile (Compound No. 98),
- 3-(4-Difluoromethoxy-3-propoxyphenyl)-5-methyl-4,5-dihydro-isoxazole-5-carbonitrile (Compound No. 99),
- 3-(4-Difluoromethoxy-3-isopropoxyphenyl)-5-methyl-4,5-dihydro-isoxazole-5-carbonitrile (Compound No. 100),
- 25 -3-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-difluoromethoxyphenyl]-5-methyl-4,5-dihydro-isoxazole-5-carbonitrile (Compound No. 101),
- 3-(3-Benzoyloxy-4-difluoromethoxyphenyl)-5-methyl-4,5-dihydro-isoxazole-5-carbonitrile (Compound No. 102),

WO 2005/051931

PCT/IB2004/003893

30

-2-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-pyridine (Compound No. 103),

-3-[3-(Cyclopentyloxy)-4-methoxyphenyl]-5-methyl-4,5-dihydroisoxazole-5-carboxylic acid hydrazide (Compound No. 104),

5 -Acetic acid (Z)-2-amino-2-{3-[3-(cyclopentyloxy)-4-methoxyphenyl]-5-methyl-4,5-dihydroisoxazol-5-yl} vinyl ester (Compound No. 105).

10





Table 1: (wherein $R_4=Y_1=Y_2=H$, $R_1=X_1=CH_3$, $X_2=cyclopentyl$, $X=Y=O$)

Compound No.	R ₂
1	
2	
3	
4	
5	
6	
7	
8	
9	


WO 2005/051931

PCT/IB2004/003893

31






10	CN
104	-CONHNH ₂
105	

5



Formula I

10 Table 2: (Formula I, wherein R₄=Y₁=Y₂=H, R₁=X₁=CH₃, X₂=cyclopentyl, X=Y=O, R₂=(CH₂)_n-NHCO-R₇, n = 1)





Compound No.	R ₇
11	
12	
13	
14	
15	

WO 2005/051931

PCT/IB2004/003893

32

Table 3: (Formula I, wherein $R_4=Y_1=Y_2=H$, $R_1=X_1=CH_3$, $X_2=cyclopentyl$, $X=Y=O$, $R_2=(CH_2)_n-NHCONHR_7$, $n = 1$)

Compound No.	R_7
16	
17	
18	
19	

5

10







Table 4: Formula XXXII (Formula I, wherein $R_4=Y_1=Y_2=H$, $R_1=X_1=CH_3$, $X_2=cyclopentyl$, $X=Y=O$, $R_2=$ )

15

Compound No.	R_2
20	
21	

WO 2005/051931

PCT/IB2004/003893














33

22	
23	
24	
25	
26	
27	
28	
29	
30	
31	
32	
33	

WO 2005/051931

PCT/IB2004/003893










34

34	
35	
36	
37	
38	
39	
40	
41	
42	
43	
44	
45	
46	

WO 2005/051931

PCT/IB2004/003893


35

47	
48	
49	
50	
51	
52	
53	
54	
55	

WO 2005/051931

PCT/IB2004/003893

36



Formula XXXIII

Table 5: Formula XXXIII (Formula I, wherein $R_4=Y_1=Y_2=H$, $R_1=X_1=CH_3$, $X_2=cyclopentyl$, $X=Y=O$, $R_2=$













[$X_7=O$])

Compound No.	R ₂
56	
57	
58	
59	
60	
61	
62	

WO 2005/051931

PCT/IB2004/003893








37

63	
64	
65	
66	
67	
68	
69	
70	
71	
72	


WO 2005/051931

PCT/IB2004/003893

38


73	
74	
75	
76	
77	
78	
79	











39



Formula XXXII

Table 6: Formula XXXII (Formula I, wherein R₄=Y₁=Y₂=H, R₁=CH₃, X=Y=O,




Compound No.	X ₂	X ₁
80	-CH ₃	-CHF ₂
81		-CHF ₂
82		-CHF ₂
83		-CHF ₂
84		-CHF ₂
85		-CHF ₂
86	CH ₃ -(CH ₂) ₃ -	-CHF ₂
87		-CHF ₂
88		-CHF ₂
89	CHF ₂	-CHF ₂
90		-CHF ₂
91		-CHF ₂
92	-CHF ₂	-CH ₃
93		-CH ₃

WO 2005/051931

PCT/IB2004/003893

40



Formula XXXIV

Table 7: Formula XXXIV (Formula I, wherein $R_4=Y_1=Y_2=H$, $R_1=CH_3$, $X=Y=O$)

Compound No.	X_1	X_2	R_{19}
94	$-CHF_2$		CN
95	$-CHF_2$		CN
96	$-CHF_2$	$-CHF_2$	CN
97	$-CH_3$		CN
98	$-CHF_2$		CN
99	$-CHF_2$		CN
100	$-CHF_2$		CN
101	$-CHF_2$		CN
102	$-CHF_2$		CN
103	$-CH_3$		

Examples set forth below demonstrate the synthetic procedures for the preparation of the representative compounds. The examples are provided to illustrate particular aspect of the disclosure and do not constrain the scope of the present invention as defined by the claims.

Experimental details

Example 1: Preparation of 3-cyclopentyloxy-4-methoxybenzaldehyde

The title compound was prepared according to methods described in *J. Med. Chem.*, (1994), 37, 1696-1703

WO 2005/051931

PCT/IB2004/003893

41

Example 2: Preparation of 3-cyclopentyloxy-4-methoxybenzaldehyde oxime

To a stirred solution of 3-cyclopentyloxy-4-methoxybenzaldehyde (0.5g, 2.2727 mmol, example 1) in ethanol (8ml) was added hydroxylamine hydrochloride (0.473g, 6.8181 mmol) and sodium acetate (0.56 g, 6.8181 mmol). The reaction mixture was allowed to stir at room temperature for 50 minutes. Ethanol was removed under reduced pressure and then residue was poured in water (20ml) and organic compound was extracted with ethyl acetate (2x15 ml). Ethyl acetate layer was dried over anhydrous sodium sulphate, filtered and finally concentrated under reduced pressure to afford compound of Formula III.

¹H NMR (CDCl₃): 9.84 (s, 1H), 8.07 (s, 1H), 6.84-7.24 (m, 3H), 4.79-4.83 (m, 1H), 3.87 (s, 3H), 1.62-2.18 (m, 8H).

Example 3: Preparation of [3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazole-5-carbonitrile (Compound No. 10)

3-cyclopentyloxy-4-methoxybenzaldehyde oxime (500 mg, 0.002 mole, example 2) was taken in 10 mL tetrahydrofuran. Methacrylonitrile (0.285 mL, 0.004 mole) was added and stirred. Sodium hypochlorite solution (10 mL, 20 times) was added dropwise. Reaction mixture was stirred vigorously at an ambient temperature. Tetrahydrofuran was removed under reduced pressure. Water was added and organic layer was extracted with ethyl acetate, dried and concentrated in vacuo. Residue was purified by column chromatography.

Yield: 63%; m.p.: 105°-106°; ¹H NMR: CDCl₃ δ= 7.33-7.34 (d, 1H), 6.96-6.99 (d, 1H), 6.84-6.87 (d, 1H), 4.80-4.84 (m, 1H), 3.86-3.88 (s, 3H), 3.80-3.86 (d, 1H), 3.36-3.41 (d, 1H), 1.80-2.0 (m, 8H), 1.56-1.63 (s, 3H); Mass (m/z) 301.5 (M⁺+1).

Example 4: Preparation of [3-(3-Cyclopentyloxy-4-methoxyphenyl)-N-hydroxy-5-methyl-4,5-dihydroisoxazole-5-carboxamidine

[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazole-5-carbonitrile (200 mg, 0.0006 mole, example 3) was dissolved in 5 mL ethanol. To it anhydrous potassium carbonate (138 mg, 0.0009 mole) and hydroxylamine hydrochloride (92 mg, 0.0013 mole) was added & refluxed. Ethanol was removed under reduced

WO 2005/051931

PCT/IB2004/003893

42

pressure, water was added. Organic layer was extracted with ethyl acetate, washed with saturated sodium chloride solution, dried and concentrated in vacuo.

Yield : 95%; Mass (m/z): 334.21 ($M^+ + 1$).

Example 5: Preparation of 3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-

5 dihydroisoxazole-5-carboxylic acid hydrazide (Compound No.104)

To the ester (300 mg, 0.00086 mole, scheme I, Formula VI), hydrazine-hydrate (0.21 mL, 0.0043 mole) was added. Reaction mixture was heated at 120°C. Reaction mixture was cooled, water was added, solid, which was separated out, was filtered and dried under vacuum.

10 Yield : 49%; m.p: 159-160°; ^1H NMR (CDCl_3): δ 8.01 (s, 1H), 7.25-7.28 (d, 1H), 6.99-7.02 (d, 1H), 6.81-6.84 (d, 1H), 4.77-4.80 (m, 1H), 3.86 (s, 3H), 3.72-3.80 (d, 1H), 3.20-3.25 (d, 1H), 1.61-2.03 (m, 11H); Mass (m/z): 334.2 ($M^+ + 1$).

Example 6: Preparation of 5-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazol-5-yl]-1H-tetrazole (Compound No. 9)

15 [3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazole-5-carbonitrile (0.00029 mole, 100 mg, example 3), sodium azide (28 mg, 0.0004 mole) and triethylamine hydrochloride (0.0005 mole, 80 mg) was taken in 20 mL toluene. Reaction mixture was refluxed overnight. Toluene was removed and then added water to it.. Extracted with ethyl acetate, washed with brine, dried and concentrated in vacuo.

20 Yield : 79%; m.p.: 161°C; ^1H NMR (MeOD): δ 7.282-7.288 (d, 1H), 7.11-7.15 (d, 1H), 6.93-6.95 (d, 1H), 4.8 (m, 1H), 3.94-4.0 (d, 1H), 3.81 (s, 3H), 3.61-3.675 (d, 1H), 1.59-1.86 (m, 11H); Mass (m/z): 344.22 ($M^+ + 1$).

Example 7: Preparation of 3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazol-5-yl]-4H-[1,2,4]oxadiazole-5-thione (Compound No. 2)

25 A mixture of [3-(3-Cyclopentyloxy-4-methoxyphenyl)-N-hydroxy-5-methyl-4,5-dihydroisoxazole-5-carboximidine (0.0006 mole, 200 mg, example 4), thiocarbonyldiimidazole (0.0009 mole, 160 mg) and 1,8-diazabicyclo[5.4.0]undec-7-one (0.002 mol-358 mL) was taken in acetonitrile and stirred at an ambient temperature. Acetonitrile was removed under reduced pressure, water was added, organic layer was

WO 2005/051931

PCT/IB2004/003893

43

extracted with ethyl acetate, washed with saturated sodium chloride solution, dried and concentrated in vacuo. The residue was purified by column chromatography.

Yield : 50%; m.p: 172°C; ¹H NMR (CDCl₃): δ 7.26 (d, 1H), 6.98-7.01 (d, 1H), 6.83-6.86 (d, 1H), 4.78-4.81 (m, 1H), 3.88-3.92 (d, 1H), 3.86 (s, 3H), 3.40-3.45 (d, 1H), 1.25-2.04 (m, 11H); Mass (m/z): 376.15 (M⁺+1).

Example 8: Preparation of 2-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 4)

To 3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazole-5-carboxylic acid hydrazide (250 mg, example 5) was added triethylorthoformate (5 mL). Reaction mixture was heated at 120°C for 3 hours. Excess triethylorthoformate was evaporated and the residue was heated at 140°C for 2 hours. Reaction mixture was diluted with water, saturated with potassium carbonate and extracted with ethyl acetate. Organic layer was dried, concentrated and purified by column chromatography.

Yield : 39%; m.p : 95°C; ¹H NMR (CDCl₃): δ 8.44 (s, 1H), 7.37 (d, 1H), 7.05-7.08 (d, 1H), 6.85-6.88 (d, 1H), 4.82-4.83 (m, 1H), 4.19-4.24 (d, 1H), 3.88 (s, 3H), 3.43-3.49 (d, 1H), 1.62-2.30 (m, 8H), 1.24-1.28 (s, 3H); Mass (m/z): 344.16 (M⁺+1).

Example 9: Preparation of 2-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-methyl-[1,3,4]oxadiazole (Compound No. 5)

Prepared as described in example 8 by using triethylortho acetate instead of triethylortho formate.

Yield : 75%; m.p : oily; ¹H NMR (CDCl₃): δ 7.364 (s, 1H), 7.04-7.07 (d, 1H), 6.84-6.87 (d, 1H), 4.816 (s, 1H), 4.16-4.21 (d, 1H), 3.88 (s, 3H), 3.37-3.43 (d, 1H), 2.556 (s, 3H), 1.621-2.15 (m, 8H), 1.25-1.31 (m, 3H); Mass (m/z) 358.23 (M⁺+1).

Example 10: Preparation of 3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazol-5-yl]-4H-[1,2,4]thiadiazol-5-one (Compound No. 3)

Amidoxime (200 mg, 0.0006 mole, scheme I, Formula VII) was taken in 3 mL tetrahydrofuran. To it thiocarbonyl diimidazole (160 mg, 0.0007 mole) was added. Reaction mixture was stirred at an ambient temperature. The reaction mixture was diluted with water, extracted with ethyl acetate, washed with water, dried and concentrated in

WO 2005/051931

PCT/IB2004/003893

44

vacuo. The residue was dissolved in tetrahydrofuran. Boron trifluoride etherate was added dropwise. The reaction mixture was stirred at an ambient temperature for 2 hours, diluted with water, extracted with ethyl acetate, dried, concentrated in vacuo and purified by column chromatography.

- 5 Yield : 23%; m.pt : 204°C; ^1H NMR (CDCl_3): δ 7.319 (s, 1H), 7.015-7.042 (d, 1H), 6.83-6.85 (d, 1H), 4.80-4.82 (m, 1H), 3.95-4.00 (d, 1H), 3.87 (s, 3H), 3.33-3.39 (d, 1H), 1.83-2.04 (m, 8H), 1.25-1.62 (m, 3H); Mass (m/z): 376.14 (M^+ +1).

Example 11: Preparation of 3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-4H-[1,2,4]oxadiazol-5-one (Compound No. 1)

- 10 Amidoxime (100 mg, 0.0003 mole, scheme I, Formula VII) was taken in dimethylformamide (1 mL). At 0°C pyridine was added then at same temperature methyl chloroformate was added dropwise. The reaction mixture was stirred at 0 °C for about 30 minutes, water was added and organic layer was extracted with ethyl acetate, washed with saturated sodium chloride solution, dried and concentrated in vacuo. To residue xylene (5
15 ml) was added and refluxed for 18 hours. Xylene was removed under reduced pressure. The crude product was purified by column chromatography.

Yield : 37%; m.pt. : oily; ^1H NMR (CDCl_3): δ 7.29 (s, 1H), 7.01-7.04 (d, 1H), 6.84-6.87 (d, 1H), 4.78-4.81 (m, 2H), 3.91-3.96 (d, 1H), 3.88 (s, 3H), 3.31-3.40 (d, 1H), 1.22-2.00 (m, 11H); Mass (m/z): 360.18 (M^+ +1).

- 20 Example 12: Preparation of 3-{3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazol-5-yl]-[1,2,4]oxadiazol-5-yl}-pyridine (Compound No. 7)

- Nicotinic acid (0.0002 mole, 30 mg) was dissolved in dry dimethylformamide (1 mL). To it, molecular sieves (100 mg, 4 Å) and triethylamine (0.0003 mole, .05 mL) was added. The reaction mixture was cooled to -20°C and isobutylchloroformate (0.0004
25 mole, .06 mL) was added. After 10 minutes amidoxime (0.0004 mole, 160 mg, scheme I, Formula VII) in dimethylformamide (2 mL) was added. The reaction mixture was stirred at an ambient temperature overnight. Some fresh molecular sieves were added. The reaction mixture was heated at 120°C for 12 hours, mixture was filtered. To filtrate water was added, extracted with ethyl acetate, washed, dried and concentrated in vacuo. The
30 residue was purified by column chromatography.

WO 2005/051931

PCT/IB2004/003893

45

Yield : 25%; m.p. : oily; ^1H NMR (CDCl_3): δ 9.38 (s, 1H), 8.83-8.84 (d, 1H), 8.42-8.44 (d, 1H), 7.47-7.51 (m, 2H), 7.06-7.09 (d, 1H), 6.85-6.87 (d, 1H), 4.81-4.83 (m, 1H), 4.07-4.13 (d, 1H), 3.88 (s, 3H), 3.41-3.46 (d, 1H), 1.62-2.09 (m, 8H), 0.8-0.98 (m, 3H); Mass (m/z): 421.40 ($\text{M}^+ + 1$).

5 The following compounds were prepared following the above procedure

—4-{3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazol-5-yl}-pyridine (Compound No. 57),

Mass (m/z): 421.40 ($\text{M}^+ + 1$)

10 —5-tert-Butyl-3-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 58),

Mass (m/z): 400.42 ($\text{M}^+ + 1$)

15 —3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-(2-ethoxy-phenyl)-[1,2,4]oxadiazole (Compound No. 59),

Mass (m/z): 464.43 ($\text{M}^+ + 1$)

m.p.: 138.5-139°C

20 —2,6-Dichloro-4-{3-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazol-5-yl}-pyridine (Compound No. 64),

Mass (m/z): 489 ($\text{M}^+ + 1$)

m.p.: 136.5 °C

25 —3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-isopropyl-[1,2,4]oxadiazole (Compound No. 65),

m.p.: 87 °C Mass (m/z): 396.00 ($\text{M}^+ + 1$)

WO 2005/051931

PCT/IB2004/003893

46

-5-Cyclohexyl-3-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 66),

Mass (m/z): 426.42 (M^+ +1)

- 5 -5-(2-Chlorophenyl)-3-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 76),

Mass (m/z): 454.31 (M^+ +1)

m.p.: 122°C

- 10 -3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-(2,4-dichloro-phenyl)-[1,2,4]oxadiazole (Compound No. 77),

Mass (m/z): 488.25 (M^+ +1)

m.p.: 129°C

- 15 Example 13: Preparation of 5-tert-Butyl-3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 8)

- Amidoxime (100 mg, 0.0003 mole, scheme I, Formula VII) was taken in benzene (2 mL). Pivaloyl chloride (0.1 mL, 0.0009 mole) was added. The reaction mixture was refluxed for about 3 hours. Benzene was removed under reduced pressure. The residue
20 was dissolved in ethyl acetate, washed with saturated sodium bicarbonate solution, dried and concentrated in vacuo. The residue was taken in dimethylformamide (5 mL) and refluxed for 3 hours. Dimethylformamide was removed under reduced pressure, water was added, extracted with ethyl acetate, dried and concentrated in vacuo.

- Yield : 25%; m.p. : sticky solid; ^1H NMR (CDCl_3): δ 7.39-7.40 (d, 1H), 7.04-7.08(d, 1H),
25 6.84-6.87 (d, 1H), 4.80-4.83 (m, 1H), 4.02-4.07 (d, 1H), 3.87 (s, 3H), 3.31-3.36 (d, 1H), 1.74-1.96 (m, 8H), 1.43 (s, 9H), 1.24-1.35 (m, 3H); Mass (m/z): 400.42 (M^+ +1).

WO 2005/051931

PCT/IB2004/003893

47

Example 14: Preparation of 3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazol-5-yl]-4-methyl-4H-[1,2,4]oxadiazole-5-thione (Compound No. 6)

3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazol-5-yl]-4H-[1,2,4]oxadiazole-5-thione (0.0001 mole, 40 mg, example 7), was dissolved in acetone (2 mL). To it potassium carbonate (0.001 mole, 147 mg) and methyl iodide (0.0002 mole, 0.016 mL) were added. The reaction mixture was refluxed for overnight. Filtered to remove potassium carbonate, washed with acetone. From filtrate, acetone was removed under reduced pressure to give a low melting solid compound.

Yield : 72%; ^1H NMR (CDCl_3): δ 7.38 (d, 1H), 7.03-7.06 (d, 1H), 6.83-6.86 (d, 1H), 4.80-4.82 (m, 1H), 3.97-4.02 (d, 1H), 3.87 (s, 3H), 3.31-3.37 (d, 1H), 2.72 (s, 3H), 1.80-1.99 (m, 8H), 1.26-1.32 (m, 3H); Mass (m/z): 390.38 ($\text{M}^+ + 1$).

Example 15: General method of preparation of compound of Formula XX (wherein $\text{R}_1 = \text{X}_1 = \text{CH}_3$, $\text{Y}_1 = \text{R}_4 = \text{Y}_2 = \text{H}$, $\text{X}_2 = \text{cyclopentyl}$ and $n = 1$)

Method A: To a stirred solution of 3-(3-Cyclopentyloxy-4-methoxy-phenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-methylamine (0.3569 mmol, 1equiv, Scheme II, Formula XIX) in 2 mL chloroform was added triethylamine (2.6767 mmol, 7.5 equiv). Compound of Formula R_{12}COCl (0.3925 mmol, 1.1equiv) was added dropwise over a period of 15 minutes with stirring the solution vigorously. The reaction was allowed to stir at an ambient temperature. The reaction mixture was quenched by adding 5 mL water. The resulting mixture was extracted with chloroform. The organic layer was thoroughly washed with water and was dried over anhydrous sodium sulphate, filtered and concentrated over buchi to afford the crude product. The crude product was purified over silica gel column (100-200 mesh) using hexane and ethyl acetate mixture as eluent.

The following compounds were prepared following the above general procedure

—Morpholine-4-carboxylic acid [3-(3-cyclopentyloxy-4-methoxy-phenyl)-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl]-amide (Compound No. 11),
Mass (m/z): 418.30 ($\text{M}^+ + 1$),

—N-[3-(3-Cyclopentyloxy-4-methoxy-phenyl)-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl]-4-fluoro-benzamide (Compound No. 12),
Mass (m/z): 427.27 ($\text{M}^+ + 1$),

WO 2005/051931

PCT/IB2004/003893

48

-Adamantane-1-carboxylic acid [3-(3-cyclopentyloxy-4-methoxy-phenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl-methyl]amide (Compound No. 13),

Mass (m/z): 467.44 ($M^+ + 1$),

5 -Furan-2-carboxylic acid [3-(3-cyclopentyloxy-4-methoxy-phenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl-methyl]-amide (Compound No. 14),

Mass (m/z): 339.24 ($M^+ + 1$),

10 -1-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl-methyl]-3-(2-trifluoromethyl)-phenyl)-urea (Compound No. 16),

Mass (m/z): 492.40 ($M^+ + 1$)

-1-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl-methyl]-3-(2,4-difluorophenyl)-urea (Compound No. 17),

15 Mass (m/z): 460.31 ($M^+ + 1$)

-1-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl-methyl]-3-O-tolyl-urea (Compound No. 18),

Mass (m/z): 438.31 ($M^+ + 1$)

20

-Morpholine-4-carboxylic acid [3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl-methyl]-amide (Compound No. 19),

Mass (m/z): 418.30 ($M^+ + 1$)

25 Method B: To a solution of 3-(3-Cyclopentyloxy-4-methoxy-phenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-methylamine (0.3407 mmol, 1 equiv, scheme II, Formula XX) and $R_{12}COOH$ (0.3407 mmol, 1 equiv.) in 0.8 mL dry dimethylformamide at 0°C was added 1-hydroxybenzotriazole (0.3407 mmol, 1 equiv) and N-methylmorpholine (1.3628 mmol, 4 equiv.). The reaction mixture was allowed to stir at 0°C for 30 minutes. Thereafter, 1-[3-(dimethylamino)propyl-3-ethyl]carbodiimide hydrochloride (0.6814 mmol, 2 equiv.) was added to the reaction mixture and reaction was continued at 0°C for 1 hour and thereafter at an ambient temperature for 20 hours. The reaction was quenched by adding water. The resulting reaction mixture was extracted with ethyl acetate. The ethyl acetate layer was dried over anhydrous sodium sulphate, concentrated in vacuo to afford the crude product.

35 The crude product was purified over silica gel column (100-200 mesh) using hexane and ethyl acetate mixture as eluent.

WO 2005/051931

PCT/IB2004/003893

49

The following compound was prepared following the above procedure (Method B)

-2-(3-Cyclopentyloxy-4-methoxy-phenyl)-N-[3-(3-cyclopentyloxy-4-methoxy-phenyl)-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl]-acetamide (Compound No. 15)

Mass (m/z): 523.37 ($M^+ + 1$)

5

Example 16: General method of preparation of compound of Formula IX (wherein $R_4=Y_1=Y_2=H$, $R_1=CH_3$, $X_1=CHF_2$, $X=Y=O$, $R_{11}=H$)

Step 1: Preparation of compound of Formula VI

Oxime (Formula IV, scheme I, 1 equiv.) and methyl methacrylate (10 equiv.) were taken in tetrahydrofuran. At ambient temperature sodium hypochlorite solution was added dropwise. The reaction mixture was stirred at room temperature for overnight. Tetrahydrofuran was removed under reduced pressure. Water was added and extracted with ethyl acetate. The mixture was washed with saturated sodium chloride solution, dried over anhydrous sodium sulfate and concentrated *in vacuo* to give pure compound.

15 Step 2: Preparation of compound of Formula VIII

To the ester compound (Formula VI, step 1) hydrazine hydrate (10 equiv.) was added and allowed to stir at 120 °C for about 3 hours. When the reaction was complete, it was cooled and water was added. Solids which separated out were filtered, dissolved in ethyl acetate, washed with water, dried over anhydrous sodium sulfate and concentrated *in vacuo* to give pure compound.

20

Step 3: Preparation of compound of Formula IX

Hydrazide (Formula VIII, step 2) and triethylorthoformate (5 ml per mmole) were heated at 120 °C for about 3 hours. Excess triethylorthoformate was evaporated and the residue heated for further about 2 hours at 140 °C. When the reaction was complete, the reaction mixture was diluted with water, saturated with potassium carbonate and extracted with ethyl acetate. Organic layer was dried and concentrated *in vacuo*. Purification was done by column chromatography to give pure compound.

25

The following compounds were prepared following the above general procedure

-2-[3-(4-Difluoromethoxy-3-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 80)

30

Mass (m/z): 326.12 ($M^+ + 1$),

WO 2005/051931

PCT/IB2004/003893

50

- 2-{3-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-difluoromethoxyphenyl]-5-methyl-4,5-dihydro-isoxazol-5-yl}-[1,3,4]oxadiazole (Compound No. 81),
Mass (m/z): (M^+ +1)
- 5

-2-{3-[3-(benzyloxy)-4-difluoromethoxyphenyl]-5-methyl-4,5-dihydro-isoxazol-5-yl}-[1,3,4]oxadiazole (Compound No. 82)
Mass (m/z): 402.11 (M^+ +1),
- 10

-2-[3-[4-Difluoromethoxy-3-ethoxyphenyl]-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 83)
Mass (m/z): 340.12 (M^+ +1),
- 15

-2-[3-[4-Difluoromethoxy-3-isopropoxyphenyl]-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 84)
Mass (m/z): 354.10 (M^+ +1),
- 20

-2-[3-(3-Cyclohexyloxy-4-difluoromethoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 85)
Mass (m/z): 394.16 (M^+ +1),
- 25

-2-[3-(3-Butoxy-4-difluoromethoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 86)
Mass (m/z): 368.09 (M^+ +1),
- 30

-2-[3-(3-Cycloheptyloxy-4-difluoromethoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 87)
Mass (m/z): 408.17 (M^+ +1),
- 35

-2-[3-(4-Difluoromethoxy-3-propoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 88)
Mass (m/z): 354.14 (M^+ +1),

-2-[3-(3,4-Bis-difluoromethoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 89)
Mass (m/z): 362.21 (M^+ +1),

-2-[3-(3-Cyclopentyloxy-4-difluoromethoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 90)
Mass (m/z): 380.19 (M^+ +1),

WO 2005/051931

PCT/IB2004/003893

51

-2-[3-(3-Cyclopropylmethoxy-4-difluoromethoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 91)

Mass (m/z): 366.18 ($M^+ + 1$),

5 -2-[3-(3-Cyclopropylmethoxy-4-methoxy-phenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 92),

Mass (m/z): 330.18 ($M^+ + 1$)

10 -2-[3-(3-Difluoromethoxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]oxadiazole (Compound No. 93)

Mass (m/z): 326.11 ($M^+ + 1$),

Example 17: General method of preparation of compound of Formula XXIV (wherein $R_4=Y_1=Y_2=H$, $R_1=X_1=CH_3$, $X_2=cyclopentyl$, $X=Y=O$)

15 Step 1: Preparation of compound of Formula XXIII

Nitriles (Formula XXII, Scheme III, 1 equiv.) was taken in solution of ethanol/water (1:4) and stirred for about 5 minutes. To this hydroxylamine hydrochloride (3.7 equiv.) and sodium carbonate (1.8 equiv.) were added and stirred for about 10 minutes at ambient temperature. The reaction mixture was stirred at reflux for about 18 hours. Ethanol was removed under reduced pressure. Water was added and triturated. Solid which precipitates out was filtered and dried under *vacuo* to give the desired amidoxime.

Step 2: Preparation of compound of Formula XXIV

25 Acid (Formula VI, Scheme I, 1 equiv.) and amidoxime (Formula XXIII, step 1, 1.1 equiv.) was taken in dry dimethylformamide. At 0 °C hydroxybenzotriazole (1 equiv.) and N-methyl morpholine (4 equiv.) were added and stirred at 0 °C for about one hour. At the same temperature 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide hydrochloride (2 equiv.) was added. The reaction mixture was stirred at room temperature for about 24 hours. Water was added, extracted with ethyl acetate, dried and concentrated *in vacuo*.
30 Solid, which formed, was taken in ethanol: water (7:1) and sodium acetate (1.5 equiv.) was added. Reaction mixture was refluxed at 80-90 °C for about 3 hours. Cooled, solid, which separated out, was filtered and recrystallized with ethanol.

The following compounds were prepared following the above general procedure

WO 2005/051931

PCT/IB2004/003893

52

–3-(2-Chloro-6-trifluoromethylphenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 20)

Mass (m/z): 522.22 ($M^+ + 1$),

- 5 –3-(2-Chloro-4-fluorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 21)

Mass (m/z): 472.22 ($M^+ + 1$),

- 10 –3-(4-Chloro-2-methoxyphenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 22)

Mass (m/z): 484.25 ($M^+ + 1$),

–3-(3-Chloro-4-fluorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 23)

- 15 Mass (m/z): 472.22 ($M^+ + 1$),

–3-(3-Chloro-4-methoxyphenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 24)

Mass (m/z): 484.25 ($M^+ + 1$),

20

–3-(3-Fluorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 25)

Mass (m/z): 438.21 ($M^+ + 1$),

- 25 –3-(3,4-Difluorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 26)

Mass (m/z): 456.2 ($M^+ + 1$),

- 30 –3-(4-Methoxyphenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 27)

Mass (m/z): 450.27 ($M^+ + 1$),

–3-(3,4-Dimethoxyphenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 28)

- 35 Mass (m/z): 480.27 ($M^+ + 1$),

–3-(2-Chloro-6-fluorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 29)

Mass (m/z): 472.22 ($M^+ + 1$),

WO 2005/051931

PCT/IB2004/003893

53

-3-(2,5-Difluoro-phenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 30)

Mass (m/z): 456.22 ($M^+ + 1$),

5 -3-(2,6-Dichlorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 31)

Mass (m/z): 488.18 ($M^+ + 1$),

10 -3-(2,3-Dichloro-phenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 32)

Mass (m/z): 488.18 ($M^+ + 1$),

-3-(2,4-Dichlorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 33)

15 Mass (m/z): 488.18 ($M^+ + 1$),

-3-(3,5-Dichlorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 34)

Mass (m/z): 488.18 ($M^+ + 1$),

20

-3-(2,5-Dichlorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 35)

Mass (m/z): 488.18 ($M^+ + 1$),

25 -3-(3,5-Difluorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 36)

Mass (m/z): 456.28 ($M^+ + 1$),

30 -3-(3-Chlorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 37)

Mass (m/z): 454.20 ($M^+ + 1$),

-3-(2,4-Difluorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 38)

35 Mass (m/z): 456.27 ($M^+ + 1$),

-3-(3,4-Dichlorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 39)

Mass (m/z): 488.21 ($M^+ + 1$),

WO 2005/051931

PCT/IB2004/003893

54

-3-(4-Chlorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 40)

Mass (m/z): 434.24 ($M^+ + 1$),

5 -4-{5-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole-3-yl}-phenylamine (Compound No. 41)

Mass (m/z): 435.27 ($M^+ + 1$),

10 -3-Phenyl-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 42)

Mass (m/z): 420.30 ($M^+ + 1$),

-3-(3,4-Dimethylphenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 43)

15 Mass (m/z): 448.32 ($M^+ + 1$),

-3-(2-Chlorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 44),

Mass (m/z): 453.5 ($M^+ + 1$),

20

-3-(4-Fluorophenyl)-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 45),

Mass (m/z): 438.29 ($M^+ + 1$),

25 -3-Methyl-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 46).

Mass (m/z): 358.0 ($M^+ + 1$),

30 Example 18: General method of preparation of compound of Formula VI (wherein $Y_1 = Y_2 = R_4 = H$, $R_1 = CH_3$, $R_r = CN$, $(CH_2)_2CN$)

Oxime (Formula IV, scheme 1, 1 equiv.) and compound of Formula V (2 equiv.) were taken in tetrahydrofuran. At ambient temperature, sodium hypochloride solution was added dropwise. The reaction mixture was stirred at room temperature overnight.

35 Tetrahydrofuran was removed under reduced pressure. Water was added, extracted with ethyl acetate, washed with saturated sodium chloride solution, dried and concentrated *in vacuo*. Purification was done by column chromatography using silica gel (100-200).

The following compounds were prepared following the above procedure

WO 2005/051931

PCT/IB2004/003893

55

—3-(3-Cyclopropylmethoxy-4-difluoromethoxyphenyl)-5-methyl-4,5-dihydro-isoxazole-5-carbonitrile (Compound No. 94),

Mass (m/z): 323.25 ($M^+ + 1$)

5 —3-(3-Cyclopentyloxy-4-difluoromethoxyphenyl)-5-methyl-4,5-dihydro-isoxazole-5-carbonitrile (Compound No. 95),

Mass (m/z): 337.14 ($M^+ + 1$)

10 —3-(3,4-Bis-difluoromethoxyphenyl)-5-methyl-4,5-dihydro-isoxazole-5-carbonitrile (Compound No. 96),

Mass (m/z): 319.15 ($M^+ + 1$)

—3-(3-Cyclopropylmethoxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazole-5-carbonitrile (Compound No. 97),

15 Mass (m/z): 287.24 ($M^+ + 1$)

—3-(3-Butoxy-4-difluoromethoxyphenyl)-5-methyl-4,5-dihydro-isoxazole-5-carbonitrile (Compound No. 98),

20 Mass (m/z): 325.13 ($M^+ + 1$)

—3-(4-Difluoromethoxy-3-propoxyphenyl)-5-methyl-4,5-dihydro-isoxazole-5-carbonitrile (Compound No. 99),

Mass (m/z): 311.07 ($M^+ + 1$)

25 —3-(4-Difluoromethoxy-3-isopropoxyphenyl)-5-methyl-4,5-dihydro-isoxazole-5-carbonitrile (Compound No. 100),

Mass (m/z): 311.15 ($M^+ + 1$)

30 —3-[3-(Bicyclo[2.2.1]hept-2-yloxy)-4-difluoromethoxyphenyl]-5-methyl-4,5-dihydro-isoxazole-5-carbonitrile (Compound No. 101),

Mass (m/z): 363.14 ($M^+ + 1$)

—3-(3-Benzylloxy-4-difluoromethoxyphenyl)-5-methyl-4,5-dihydro-isoxazole-5-carbonitrile (Compound No. 102),

35 Mass (m/z): 360.18 ($M^+ + 1$)

WO 2005/051931

PCT/IB2004/003893

56

Example 19: Preparation of 3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-methyl-[1,2,4]oxadiazole (Compound No. 56)

To a mixture of amidoxime (100 mg, 0.00030 mole, Scheme I, Formula VII) and powdered molecular sieves (4A°, 500 mg), dry tetrahydrofuran (3 ml) was added. The reaction mixture was stirred for about 30 minutes. Sodium hydride (11 mg, 0.0003 mole) was added and heated at about 60°C for about 45 minutes. Methyl acetate (0.047 ml, 0.0006 mole) was added to reaction mixture and refluxed at about 65-70°C for one hour. The reaction mixture was filtered, and washed with ethyl acetate. The organic layer was concentrated under reduced pressure to give crude product, which was purified by column chromatography using silica gel (100-200). Yield: 50 mg.

The following compounds were prepared following the above procedure

-3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-fluoromethyl-[1,2,4]oxadiazole (Compound No. 67),

Mass (m/z): 376.22 ($M^+ + 1$)

-3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-(3-fluorophenyl)-[1,2,4]oxadiazole (Compound No. 70),

Mass (m/z): 438.24 ($M^+ + 1$)

-{3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazol-5-yl}-acetonitrile (Compound No. 71),

Mass (m/z): 383.24 ($M^+ + 1$)

Example 20: Preparation of 3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-(tetrahydro-furan-2-yl)-[1,2,4]oxadiazole (Compound No. 68)

Amidoxime (100 mg, 0.0003 mole, Scheme I, Formula VII) and tetrahydro-2-furoic acid (0.03 ml, 0.0003 mole) was taken in dimethylformamide (1 ml). At about 0°C, 1-hydroxybenzotriazole (40 mg, 0.0003 mole) and N-methylmorpholine (0.168 ml, 0.0012 mole) were added and stirred for about 1 hour. Thereafter, at about 0°C 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide hydrochloride (115 mg, 0.0006 mole) was added and stirred at room temperature for about 24 hours. Water (5 ml) was added, extracted with ethyl acetate. The ethyl acetate layer was dried over anhydrous sodium

WO 2005/051931

PCT/IB2004/003893

57

sulphate, and concentrated *in vacuo*. Dimethylformamide (2 ml) was added to the reaction residue. 50 mg powdered molecular sieves was added and refluxed at about 110-120°C for about 4 hours. The resultant was filtered and washed with ethyl acetate. The organic layer was concentrated and purified by column chromatography using silica gel (100-200).

5 Yield: 40 mg

The following compounds were prepared similarly

–3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-cyclopropyl-[1,2,4]oxadiazole (Compound No. 60),

Mass (m/z): 384.31 ($M^+ + 1$); m.p.: 105-106°C.

10

–5-(3-Chlorophenyl)-3-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 61),

Mass (m/z): 454.31 ($M^+ + 1$); m.p.: 103-104°C.

15

–5-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-m-tolyl-[1,2,4]oxadiazole (Compound No. 62),

Mass (m/z): 434.42 ($M^+ + 1$); m.p.: 131-132°C.

20

–3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-(3,5-dimethyl-phenyl)-[1,2,4]oxadiazole (Compound No. 63),

Mass (m/z): 448.00 ($M^+ + 1$); m.p.: 131°C.

25

–3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-(4-fluoro-phenyl)-[1,2,4]oxadiazole (Compound No. 69),

Mass (m/z): 438.26 ($M^+ + 1$); m.p.: 146.1-146.3°C.

30

–4-{3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole-5-yl}-benzonitrile (Compound No. 72),

Mass (m/z): 445.32 ($M^+ + 1$)

–3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-trifluoromethyl-[1,2,4]oxadiazole (Compound No. 73),

Mass (m/z): 412.25 ($M^+ + 1$)

35

–3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-(3-methoxy-phenyl)-[1,2,4]oxadiazole (Compound No. 74),

WO 2005/051931

PCT/IB2004/003893

58

Mass (m/z): 450.26 ($M^+ + 1$); m.p.: 121-122°C.

-3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-(3,4-dimethoxy-phenyl)-[1,2,4]oxadiazole (Compound No. 75),

5 Mass (m/z): 480.32 ($M^+ + 1$); m.p.: 119-120°C.

-5-Cyclopentyl-3-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,2,4]oxadiazole (Compound No. 78),

Mass (m/z): 412.33 ($M^+ + 1$)

10

-3-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-5-(3,4-dichloro-phenyl)-[1,2,4]oxadiazole (Compound No. 79),

Mass (m/z): 488.17 ($M^+ + 1$); m.p.: 113.5-114.9°C.

15 Example 21: Preparation of 2-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-pyridine (Compound No. 103)

3-cyclopentyloxy-4-methoxybenzaldehyde oxime (250 mg, 1.063 mmol) and 2-vinyl-pyridine (167 mg, 1.595 mmol) were taken in tetrahydrofuran (3 ml). The reaction mixture was stirred for about 10 minutes. Sodium hypochloride solution (1 ml, 10.63 mmol) was added gradually over 15 minutes and stirred for 2 hours. THF was evaporated off and the residue was extracted with ethylacetate. The organic layer was washed with water, dried over anhydrous sodium sulphate and concentrated. The product was purified using column chromatography. Mass (m/z): 339.21 ($M^+ + 1$).

25 Example 22: Preparation of {5-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4] thiadiazol-2-yl}-cyclopropylamine (Compound No. 49)

3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazole-5-carboxylic acid (250 mg, 0.00078 mole) and cyclopropylthiosemicarbazide (102 mg, 0.00078 mole) were taken in dioxane (10 ml). At about 65°C, POCl₃ (0.07 ml, 0.00078) was added to the reaction mixture. The reaction mixture was refluxed at about 65°C for about 5 hours and then at room temperature overnight. Dioxane was removed under reduced pressure. Saturated sodium bicarbonate solution was added. Extraction was done by ethyl acetate, and the extract was washed with saturated sodium chloride solution. The product was dried over anhydrous sodium sulphate and concentrated *in vacuo* to give solid

30

WO 2005/051931

PCT/IB2004/003893

59

compound, which was further crystallized by ethanol to give white solid compound having m.pt- 188-189°C. Yield = 44 mg; Mass ($M^+ + 1$) = 415.

The following compound was prepared similarly

5 -5-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-[1,3,4]thiadiazol-2-yl-amine (Compound No. 48)

Mass (m/z): 375.37 ($M^+ + 1$)

Example 23: Preparation of 3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-(4,5-dihydro-thiazol-2-yl)-5-methyl-4,5-dihydro-isoxazole (Compound No. 51)

10 3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydroisoxazole-5-carbonitrile (70 mg, 0.0002 mole) and 2-aminoethanthiol hydrochloride (53 mg, 0.0004 mole) were taken in 5 ml ethanol. Triethylamine (0.04 ml, 0.0003 ml) was added to the reaction mixture and refluxed for about 5 hours. Ethanol was removed under reduced pressure to get crude compound, which was purified by column chromatography using
15 silica gel (100-200). Yield: 50 mg; m.pt.: sticky solid.

Example 24: Preparation of 1-{5-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-2-ethyl-2-methyl-[1,3,4]oxadiazol-3-yl}-ethanone (Compound No. 50)

20 Step a: Hydrazide (100 mg, 0.0003 mole, Scheme IA, Formula VIII) was taken in ethanol (5 ml). Ethylmethyl ketone (0.03 ml, 0.0004 mole) was added. The reaction mixture was stirred at refluxing temperature for about 10 hours. Ethane was removed under reduced pressure to give oily compound.

25 Step b: The compound from step a was taken in pyridine (3 ml). One ml acetic anhydride was added and stirred at about 100°C for about 8 hours. A mixture of acetic anhydride and pyridine was removed under reduced pressure. 5 ml cold water was added and extraction was done by ethyl acetate. The resultant was washed with saturated sodium chloride solution, dried over anhydrous sodium sulphate and concentrated *in vacuo* to give crude compound, which was purified by column chromatography using silica gel (100-200).

WO 2005/051931

PCT/IB2004/003893

60

Example 25: Preparation of 3'-(3-Cyclopentyloxy-4-methoxyphenyl)-5,5'-dimethyl-4,5,4',5'-tetrahydro-[3,5'] biisoxazolyl-5-carbonitrile (Compound No. 52)

Step a: Oxime (350 mg, 0.00148 mole, Scheme I, Formula IV) and methacrolein (0.73 ml, 0.0089 mole) was taken in tetrahydrofuran (10 ml). Sodium hypochlorite solution (10 ml) was added dropwise to residue mixture. The reaction mixture was stirred at room temperature for about 14-16 hours. Tetrahydrofuran was removed under reduced pressure. Water (10 ml) was added, extracted with ethyl acetate, washed with saturated sodium chloride solution, dried over anhydrous sodium sulphate and concentrated *in vacuo* to give oily compound.

Step b: The compound from step a was taken in ethanol (10 ml) and to it hydroxylamine hydrochloride (160 mg, 0.0023 mole) and anhydrous sodium acetate (189 mg, 0.0028 mole) were added. The reaction mixture was stirred at room temperature for about one and half an hours. Ethanol was removed under reduced pressure, water was added, extracted with ethyl acetate. Dried over anhydrous sodium sulphate and concentrated *in vacuo* to give oily compound.

Step c:- The compound from step b was taken in tetrahydrofuran (5 ml) and to it methacrylonitrile (0.246 ml, 0.0036 mole) was added. Sodium hypochlorite solution (3 ml) was added to reaction mixture dropwise within a 15 minute interval. The reaction mixture was stirred for about 15-16 hours. Tetrahydrofuran was removed under reduced pressure. Water (30 ml) was added, extracted with ethyl acetate. Dried over anhydrous sodium sulphate and concentrated *in vacuo* to give oily compound, which was purified by column chromatography using silica gel (100-200). Yield: 50 mg; m.pt: oily.

Example 26: Preparation of 5-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-3H-[1,3,4]oxadiazole-2-thione (Compound No. 55)

Hydrazide (70 mg, 0.0002 mole, Scheme IA, Formula VIII) was taken in ethanol (5 ml). To it potassium hydroxide solution (0.11 g, 0.0002 mole) in 1 ml ethanol were added followed by carbon disulfide (1 ml). The reaction mixture was refluxed for about 8 hours. Ethanol was removed under reduced pressure. The reaction mixture was neutralized by dilute hydrochloride (2N), and extracted with ethyl acetate. The resultant was washed with saturated sodium chloride solution, dried over anhydrous sodium

WO 2005/051931

PCT/IB2004/003893

61

sulphate and concentrated *in vacuo* to give crude product, which was purified by column chromatography using silica gel 100-200. Yield: 70 mg; m.pt: 195.5-200°C.

Example 27: Preparation of 4-Amino-5-[3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-4H-[1,2,4]triazole-3-thiol (Compound No. 54)

5 A mixture of 5-[3-(3-Cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazol-5-yl]-3H-[1,3,4]oxadiazole-2-thione (50 mg, 0.00013 mole, Example 26) and hydrazine hydrate (0.02 ml, 0.0003 mole) in ethanol (2 ml) were refluxed for about 6 hours. The solvent and excess hydrazine hydrate were removed under reduced pressure. Water was added, and the aqueous phase was extracted with ethyl acetate. The extract
10 was washed with brine, dried over anhydrous sodium sulphate and concentrated *in vacuo* to give crude product, which was recrystallized using ethyl acetate hexane (20:80). Yield: 15 mg; m.pt.: 228-229°C.

Example 28: Preparation of Acetic acid (Z)-2-amino-2-{3-[3-(cyclopentyloxy)-4-methoxyphenyl]-5-methyl-4,5-dihydroisoxazol-5-yl} vinyl ester (Compound No. 105)

15 [3-(3-Cyclopentyloxy-4-methoxyphenyl)-N-hydroxy-5-methyl-4,5-dihydroisoxazole-5-carboxamidine (Example 4, 0.0007 mole, 250 mg) was dissolved in dichloromethane. To it acetic anhydride (0.0007 mole, 0.07 ml) and triethyl amine (0.0007 mole, 0.105 ml) were added. The reaction mixture was stirred at an ambient temperature for about 2 hours. The mixture was washed with water. The organic layer
20 was dried over anhydrous sodium sulphate, concentrated *in vacuo* and the residue was purified over column chromatography. Yield: 46%; m.pt.: 130.9°C; ¹H NMR (CDCl₃): δ 7.29-7.30 (d, 1H), 7.036-7.06 (d, 1H), 6.83-6.86 (d, 1H), 5.259 (s, 2H), 4.78-4.81 (m, 1H), 3.98 (d, 1H), 3.87 (s, 3H), 3.276-3.33 (d, 1H), 2.16(s, 3H), 1.79-2.09(m, 8H), 1.25-1.29(m, 3H); Mass (m/z): 376.24 (M⁺+1).

25 Example 29: Preparation of 3-{3-[3-(cyclopentyloxy)-4-methoxyphenyl]-5-methyl-4,5-dihydroisoxazol-5-yl}-5-methyl-1,4,2-dioxazole-5-carboxylic acid ethyl ester (Compound No. 53)

To a solution of 3-(3-cyclopentyloxy-4-methoxyphenyl)-5-methyl-4,5-dihydro-isoxazole-5-carbaldehyde oxime (0.480g, 1.5116 mmole) and 2-oxo propionic acid ethyl
30 ester (1.052g, 9.0698 mmole) in tetrahydrofuran was added bleach over a period of about

WO 2005/051931

PCT/IB2004/003893

62

40 minutes. The reaction mixture was allowed to stir at room temperature for about one and half hours. Thereafter, tetrahydrofuran was removed over buchi. To the residue was added water (20 mL) and the resulting solution was extracted with ethyl acetate. Ethyl acetate layer was dried over anhydrous sodium sulphate and finally concentrated to afford
5 an oily residue, which was purified by column chromatography. Yield: 0.200g; m.p: 139-140 °C; Mass (m/z): 376.24 (M^+ +1).

Example 30: Efficacy of compounds as PDE IV inhibitors

PDE-IV Enzyme Assay

The efficacy of compounds of PDE-4 inhibitors was determined by an enzyme
10 assay using U937 cell cytosolic fraction (*BBRC*, 197: 1126-1131, 1993). Hydrolysis of cAMP to AMP was monitored using HPLC and [^3H]cAMP in the sample was detected using FLO-ONE Detector.

The enzyme preparation was incubated in the presence and absence of the test
compound for 30 min and amount/ [^3H]cAMP measured in the sample. The IC_{50} values
15 were found to be in the range of double-digit nM to >10 μM concentration.


WO 2005/051931

PCT/IB2004/003893

63

We Claim:

1. Compounds having the structure of Formula I:



Formula I

their pharmaceutically acceptable salts, pharmaceutically acceptable solvates, enantiomers, diastereomers or N-oxides wherein

1) when X is oxygen in Formula I:

R₁ is selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl; cyano; nitro; amino; substituted amino; hydroxyl; alkoxy; aryloxy; COR'; COOR'

(wherein R' can be hydrogen, alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heterocyclyl, (heterocyclyl)alkyl, or (heteroaryl)alkyl);

aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl) alkyl; (heterocyclyl) alkyl; (CH₂)₁₋₄OR'

(wherein R' is as defined above, but also including hydroxy); C(=O)NR_xR_y

(wherein R_x and R_y can be independently selected from hydrogen, alkyl, C₃₋₆

alkenyl, C₃₋₆ alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heteroaryl,

heterocyclyl, heteroarylalkyl, or heterocyclylalkyl); or

(CH₂)_m-C(=O)R₃

[wherein m is an integer in the range of 0-2 and R₃ can be optionally substituted R_p or R_q (wherein R_p can be a 4-12 membered (un)saturated monocyclic or bicyclic ring containing 1-4 heteroatom(s) selected from N, O and S wherein the ring can be attached to (CH₂)_mC(=O) through N and R_q can be a 4-12 membered (un)saturated monocyclic or bicyclic ring containing 0-4 heteroatom(s) selected from the group consisting of N, O and S wherein the ring can be attached to (CH₂)_mC(=O) through C) and wherein the substituents of R₃ can be one or more of: alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl, halogen, hydroxyl, alkoxy, aryloxy, nitro, cyano, amino, substituted amino, hydroxyalkyl, oxo, acyl,

WO 2005/051931

PCT/IB2004/003893

64

- 31 optionally substituted amino (wherein the substituents are selected from C₁-C₆
 32 alkyl, aryl, aralkyl, or cycloalkyl), aryl, carboxyl, alkaryl, carbamoyl, alkyl ether,
 33 C(=O)NR₅R₆ (wherein R₅ and R₆ are independently selected from hydrogen,
 34 alkyl, C₃₋₆ alkenyl, C₃₋₆ alkynyl, aryl, and aralkyl), optionally substituted
 35 monocyclic or bicyclic 4-12 membered carbocyclic ring system (wherein the
 36 optional substituent(s) is/are selected from alkyl, alkenyl, alkynyl, halogen,
 37 hydroxyl, and alkoxy), heteroaryl, heterocyclyl, heteroarylalkyl, or
 38 heterocyclylalkyl];
- 39 R₂ is selected from: cyano; heteroaryl; heterocyclyl; or (CH₂)_nNHCOR₇ (wherein n
 40 represents an integer 1 to 6 and R₇ can represent hydrogen, alkyl, alkenyl, alkynyl,
 41 (un)saturated, cycloalkyl, alkoxy, aryloxy, aryl, aralkyl, heteroaryl, heterocyclyl,
 42 (CH₂)₁₋₄OR' wherein R' is the same as defined above, or NR_xR_y wherein R_x and R_y are the
 43 same as defined above);
- 44 R₄ is selected from: hydrogen; alkyl; halogen; cyano; carboxy; or C(=O)NR_xR_y wherein
 45 R_x and R_y are the same as defined above;
- 46 X₁ and X₂ are independently selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl;
 47 acyl; aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl)alkyl; or (heterocyclyl)alkyl;
- 48 Y is selected from: an oxygen atom; a sulphur atom; or NR
 49 (wherein R is selected from hydrogen, alkyl, alkenyl, alkynyl, un(saturated)
 50 cycloalkyl, acyl, aryl, aralkyl, heteroaryl, heterocyclyl, (heteroaryl)alkyl, or
 51 (heterocyclyl)alkyl);
- 52 Y₁ and Y₂ are independently selected from: hydrogen; alkyl; nitro; cyano; halogen; OR
 53 wherein R is the same as defined earlier; SR wherein R is the same as defined earlier;
 54 NHR wherein R is the same as defined earlier; COOR'; or COR' wherein R' is the same
 55 as defined above, or further, Y₁ and X₂, X₁ and Y₂, X₁ and X₂ may together form a ring
 56 fused with the ring A containing 3-5 carbon atoms within the ring and having 1-3
 57 heteroatoms selected from N, O or S; and
- 58 2) when X is NR₈ or S wherein R₈ is hydrogen, lower alkyl (C₁-C₆) or aryl;
- 59 R₁, R₄, X₁, X₂, Y, Y₁ and Y₂ are the same as defined above;


WO 2005/051931

PCT/IB2004/003893

65

60 R_2 is selected from: $(CH)_nNHCOR_7$ (wherein n represents an integer 1 to 6 and R_7 is the
61 same as defined above),
62 with the proviso that when R_2 is heterocyclyl, R_1 can not be $(CH_2)_{1-4}OR'$, $C(=O)NR_xR_y$ or
63 $(CH_2)_m-C(=O)R_3$.

1 2. A compound having the structure of Formula XXXIV,



Formula XXXIV

8 their pharmaceutically acceptable salts, pharmaceutically acceptable solvates,
9 enantiomers, diastereomers or N-oxides
10 wherein

11 R_1 is selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl; cyano; nitro; amino;
12 substituted amino; hydroxyl; alkoxy; aryloxy; COR' ; $COOR'$
13 (wherein R' can be hydrogen, alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl,
14 aryl, aralkyl, heterocyclyl, (heterocyclyl)alkyl, or (heteroaryl)alkyl);
15 aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl) alkyl; (heterocyclyl) alkyl; $(CH_2)_{1-4}OR'$
16 (wherein R' is as defined above, but also including hydroxy); $C(=O)NR_xR_y$
17 (wherein R_x and R_y can be independently selected from hydrogen, alkyl, C_{3-6}
18 alkenyl, C_{3-6} alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heteroaryl,
19 heterocyclyl, heteroarylalkyl, or heterocyclylalkyl); or
20 $(CH_2)_m-C(=O)R_3$

21 [wherein m is an integer in the range of 0-2 and R_3 can be optionally substituted
22 R_p or R_q (wherein R_p can be a 4-12 membered (un)saturated monocyclic or
23 bicyclic ring containing 1-4 heteroatom(s) selected from N, O and S wherein the
24 ring can be attached to $(CH_2)_mC(=O)$ through N and R_q can be a 4-12 membered
25 (un)saturated monocyclic or bicyclic ring containing 0-4 heteroatom(s) selected
26 from the group consisting of N, O and S wherein the ring can be attached to

WO 2005/051931

PCT/IB2004/003893

66

27 (CH₂)_mC(=O) through C) and wherein the substituents of R₃ can be one or more
 28 of: alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl, halogen, hydroxyl, alkoxy,
 29 aryloxy, nitro, cyano, amino, substituted amino, hydroxyalkyl, oxo, acyl,
 30 optionally substituted amino (wherein the substituents are selected from C₁-C₆
 31 alkyl, aryl, aralkyl, or cycloalkyl), aryl, carboxyl, alkaryl, carbamoyl, alkyl ether,
 32 C(=O)NR₅R₆ (wherein R₅ and R₆ are independently selected from hydrogen,
 33 alkyl, C₃₋₆ alkenyl, C₃₋₆ alkynyl, aryl, and aralkyl), optionally substituted
 34 monocyclic or bicyclic 4-12 membered carbocyclic ring system (wherein the
 35 optional substituent(s) is/are selected from alkyl, alkenyl, alkynyl, halogen,
 36 hydroxyl, and alkoxy), heteroaryl, heterocyclyl, heteroarylalkyl, or
 37 heterocyclylalkyl];

38 R₄ is selected from: hydrogen; alkyl; halogen; cyano; carboxy; or C(=O)NR_xR_y wherein
 39 R_x and R_y are the same as defined above;

40 X₁ and X₂ are independently selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl;
 41 acyl; aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl)alkyl; or (heterocyclyl)alkyl;

42 Y is selected from: an oxygen atom; a sulphur atom; or NR
 43 (wherein R is selected from hydrogen, alkyl, alkenyl, alkynyl, un(saturated)
 44 cycloalkyl, acyl, aryl, aralkyl, heteroaryl, heterocyclyl, (heteroaryl)alkyl, or
 45 (heterocyclyl)alkyl);

46 Y₁ and Y₂ are independently selected from: hydrogen; alkyl; nitro; cyano; halogen; OR
 47 wherein R is the same as defined earlier; SR wherein R is the same as defined earlier;
 48 NHR wherein R is the same as defined earlier; COOR'; or COR' wherein R' is the same
 49 as defined above, or further, Y₁ and X₂, X₁ and Y₂, X₁ and X₂ may together form a ring
 50 fused with the ring A containing 3-5 carbon atoms within the ring and having 1-3
 51 heteroatoms selected from N, O or S; and

52 R₁₉ represents -CONHNH₂, or


53
$$\begin{array}{c} \text{---C=N---O---C(=O)---R'} \\ | \\ \text{NH}_2 \end{array}$$
, wherein R' is the same as defined for Formula I.

WO 2005/051931

PCT/IB2004/003893

67

54 3. The compound of claim 1 having the structure of Formula XXXII,



Formula XXXII

62 their pharmaceutically acceptable salts, pharmaceutically acceptable solvates,
63 enantiomers, diastereomers or N-oxides wherein

64 wherein

65 R₁ is selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl; cyano; nitro; amino;
66 substituted amino; hydroxyl; alkoxy; aryloxy; COR'; COOR'

67 (wherein R' can be hydrogen, alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl,
68 aryl, aralkyl, heterocyclyl, (heterocyclyl)alkyl, or (heteroaryl)alkyl);
69 aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl) alkyl; (heterocyclyl) alkyl; (CH₂)₁₋₄OR'
70 (wherein R' is as defined above, but also including hydroxy); C(=O)NR_xR_y

71 (wherein R_x and R_y can be independently selected from hydrogen, alkyl, C₃₋₆
72 alkenyl, C₃₋₆ alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heteroaryl,
73 heterocyclyl, heteroarylalkyl, or heterocyclylalkyl); or

74 (CH₂)_m-C(=O)R₃

75 [wherein m is an integer in the range of 0-2 and R₃ can be optionally substituted
76 R_p or R_q (wherein R_p can be a 4-12 membered (un)saturated monocyclic or
77 bicyclic ring containing 1-4 heteroatom(s) selected from N, O and S wherein the
78 ring can be attached to (CH₂)_mC(=O) through N and R_q can be a 4-12 membered
79 (un)saturated monocyclic or bicyclic ring containing 0-4 heteroatom(s) selected
80 from the group consisting of N, O and S wherein the ring can be attached to
81 (CH₂)_mC(=O) through C) and wherein the substituents of R₃ can be one or more
82 of: alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl, halogen, hydroxyl, alkoxy,
83 aryloxy, nitro, cyano, amino, substituted amino, hydroxyalkyl, oxo, acyl,

WO 2005/051931

PCT/IB2004/003893

68

84 optionally substituted amino (wherein the substituents are selected from C₁-C₆
85 alkyl, aryl, aralkyl, or cycloalkyl), aryl, carboxyl, alkaryl, carbamoyl, alkyl ether,
86 C(=O)NR₅R₆ (wherein R₅ and R₆ are independently selected from hydrogen,
87 alkyl, C₃₋₆ alkenyl, C₃₋₆ alkynyl, aryl, and aralkyl), optionally substituted
88 monocyclic or bicyclic 4-12 membered carbocyclic ring system (wherein the
89 optional substituent(s) is/are selected from alkyl, alkenyl, alkynyl, halogen,
90 hydroxyl, and alkoxy), heteroaryl, heterocyclyl, heteroarylalkyl, or
91 heterocyclylalkyl];

92 R₄ is selected from: hydrogen; alkyl; halogen; cyano; carboxy; or C(=O)NR_xR_y wherein
93 R_x and R_y are the same as defined above;

94 Y is selected from: an oxygen atom; a sulphur atom; or NR
95 (wherein R is selected from hydrogen, alkyl, alkenyl, alkynyl, un(saturated)
96 cycloalkyl, acyl, aryl, aralkyl, heteroaryl, heterocyclyl, (heteroaryl)alkyl, or
97 (heterocyclyl)alkyl);

98 Y₁ and Y₂ are independently selected from: hydrogen; alkyl; nitro; cyano; halogen; OR
99 wherein R is the same as defined earlier; SR wherein R is the same as defined earlier;
100 NHR wherein R is the same as defined earlier; COOR'; or COR' wherein R' is the same
101 as defined above, or further, Y₁ and X₂, X₁ and Y₂, X₁ and X₂ may together form a ring
102 fused with the ring A containing 3-5 carbon atoms within the ring and having 1-3
103 heteroatoms selected from N, O or S;

104 X₁ represents alkyl;

105 X₂ represents alkyl, cycloalkyl or aralkyl;


106 X₃, X₄, X₅ and X₆ independently represent C, CH, CH₂, CO, CS, NH, N, O, S; R₁₅,

107 R₁₆, and R₁₇ independently represent no atom, alkyl, COCH₃, COOC₂H₅, NH₂,

108 NH-cyclopropyl, CN, SH; and

109 --- represents an optional single bond.

1 4. The compound of claim 1 having the structure of Formula XXIII,



10 their pharmaceutically acceptable salts, pharmaceutically acceptable solvates,
11 enantiomers, diastereomers or N-oxides wherein

12 wherein

13 R₁ is selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl; cyano; nitro; amino;
14 substituted amino; hydroxyl; alkoxy; aryloxy; COR'; COOR'

15 (wherein R' can be hydrogen, alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl,
16 aryl, aralkyl, heterocyclyl, (heterocyclyl)alkyl, or (heteroaryl)alkyl);

17 aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl) alkyl; (heterocyclyl) alkyl; (CH₂)₁₋₄OR'

18 (wherein R' is as defined above, but also including hydroxy); C(=O)NR_xR_y

19 (wherein R_x and R_y can be independently selected from hydrogen, alkyl, C₃₋₆
20 alkenyl, C₃₋₆ alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heteroaryl,
21 heterocyclyl, heteroarylalkyl, or heterocyclylalkyl); or

22 (CH₂)_m-C(=O)R₃

23 [wherein m is an integer in the range of 0-2 and R₃ can be optionally substituted
24 R_p or R_q (wherein R_p can be a 4-12 membered (un)saturated monocyclic or
25 bicyclic ring containing 1-4 heteroatom(s) selected from N, O and S wherein the
26 ring can be attached to (CH₂)_mC(=O) through N and R_q can be a 4-12 membered
27 (un)saturated monocyclic or bicyclic ring containing 0-4 heteroatom(s) selected
28 from the group consisting of N, O and S wherein the ring can be attached to
29 (CH₂)_mC(=O) through C) and wherein the substituents of R₃ can be one or more
30 of: alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl, halogen, hydroxyl, alkoxy,
31 aryloxy, nitro, cyano, amino, substituted amino, hydroxyalkyl, oxo, acyl,

WO 2005/051931

PCT/IB2004/003893

70

32 optionally substituted amino (wherein the substituents are selected from C₁-C₆
33 alkyl, aryl, aralkyl, or cycloalkyl), aryl, carboxyl, alkaryl, carbamoyl, alkyl ether,
34 C(=O)NR₅R₆ (wherein R₅ and R₆ are independently selected from hydrogen,
35 alkyl, C₃₋₆ alkenyl, C₃₋₆ alkynyl, aryl, and aralkyl), optionally substituted
36 monocyclic or bicyclic 4-12 membered carbocyclic ring system (wherein the
37 optional substituent(s) is/are selected from alkyl, alkenyl, alkynyl, halogen,
38 hydroxyl, and alkoxy), heteroaryl, heterocyclyl, heteroarylalkyl, or
39 heterocyclylalkyl];

40 R₄ is selected from: hydrogen; alkyl; halogen; cyano; carboxy; or C(=O)NR_xR_y wherein
41 R_x and R_y are the same as defined above;

42 X₁ and X₂ are independently selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl;
43 acyl; aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl)alkyl; or (heterocyclyl)alkyl;

44 Y is selected from: an oxygen atom; a sulphur atom; or NR
45 (wherein R is selected from hydrogen, alkyl, alkenyl, alkynyl, un(saturated)
46 cycloalkyl, acyl, aryl, aralkyl, heteroaryl, heterocyclyl, (heteroaryl)alkyl, or
47 (heterocyclyl)alkyl);

48 Y₁ and Y₂ are independently selected from: hydrogen; alkyl; nitro; cyano; halogen; OR
49 wherein R is the same as defined earlier; SR wherein R is the same as defined earlier;
50 NHR wherein R is the same as defined earlier; COOR'; or COR' wherein R' is the same
51 as defined above, or further, Y₁ and X₂, X₁ and Y₂, X₁ and X₂ may together form a ring
52 fused with the ring A containing 3-5 carbon atoms within the ring and having 1-3
53 heteroatoms selected from N, O or S;

54 X₇ represents O or S; and

55 R₁₈ represents hydrogen, alkyl, aryl, heteroaryl, cycloalkyl or heterocyclyl.

1 5. The compound of claim 1 wherein R₂ is cyano.

1 6. The compound of claim 1 wherein R₂ is (CH₂)_nNHCOR₇, n represents an integer 1
2 to 6; and R₇ can represent hydrogen, alkyl, alkenyl, alkynyl, (un)saturated, cycloalkyl,
3 alkoxy, aryloxy, aryl, aralkyl, heteroaryl, heterocyclyl, (CH₂)₁₋₄OR' wherein R' is the same
4 as defined above, or NR_xR_y (wherein R_x and R_y can be independently selected from

WO 2005/051931

PCT/IB2004/003893

71


- 5 hydrogen, alkyl, C₃₋₆ alkenyl, C₃₋₆ alkynyl, (un)saturated cycloalkyl, aryl, aralkyl,
 - 6 heteroaryl, heterocyclyl, heteroarylalkyl, or heterocyclylalkyl).
- 1 7. The compound of claim 1 wherein R₂ is 6-membered heteroaryl.
 - 1 8. A pharmaceutical composition comprising a therapeutically effective amount of a
2 compound of claim 1, together with at least one pharmaceutically acceptable
3 carrier, excipient or diluent.
 - 1 9. A method for treating, preventing, inhibiting or suppressing an inflammatory
2 condition or disease in a patient, comprising administering to the said patient a
3 therapeutically effective amount of a compound of claim 1.
 - 1 10. A method for treating, preventing, inhibiting or suppressing an inflammatory
2 condition or disease in a patient, comprising administering to the said patient a
3 therapeutically effective amount of a pharmaceutical composition of claim 8.
 - 1 11. A method for the treatment, prevention, inhibition or suppression of AIDS, asthma,
2 arthritis, bronchitis, chronic obstructive pulmonary disease (COPD), psoriasis,
3 allergic rhinitis, shock, atopic dermatitis, crohn's disease, adult respiratory distress
4 syndrome (ARDS), eosinophilic granuloma, allergic conjunctivitis, osteoarthritis,
5 ulcerative colitis and other inflammatory diseases in a patient comprising
6 administering to said patient a therapeutically effective amount of a compound of
7 claim 1.
 - 1 12. A method for the treatment, prevention, inhibition or suppression of AIDS, asthma,
2 arthritis, bronchitis, chronic obstructive pulmonary disease (COPD), psoriasis,
3 allergic rhinitis, shock, atopic dermatitis, crohn's disease, adult respiratory distress
4 syndrome (ARDS), eosinophilic granuloma, allergic conjunctivitis, osteoarthritis,
5 ulcerative colitis and other inflammatory diseases in a patient comprising
6 administering to said patient a therapeutically effective amount of a pharmaceutical
7 composition of claim 8.

WO 2005/051931

PCT/IB2004/003893

72


1 13. A method for the preparation of compounds of Formula VII (a),



Formula VII(a)


7 their pharmaceutically acceptable salts, pharmaceutically acceptable solvates,
8 enantiomers, diastereomers or N-oxides, the method comprising:

9 reacting a compound of Formula II



Formula II

15 with a compound of Formula X₂Z (wherein Z is halogen) to give a compound of Formula
16 III, wherein



Formula III

21 X₁ and X₂ are independently selected from: alkyl; alkenyl; alkynyl; cycloalkyl; acyl; aryl;
22 aralkyl; heteroaryl; heterocyclyl; (heteroaryl)alkyl; or (heterocyclyl)alkyl;
23 Y₁ and Y₂ are independently selected from: hydrogen; alkyl; nitro; cyano; halogen; OR
24 wherein R is the same as defined earlier; SR wherein R is the same as defined earlier;
25 NHR wherein R is the same as defined earlier; COOR'; or COR' wherein R' is the same
26 as defined above, or further, Y₁ and X₂, X₁ and Y₂, X₁ and X₂ may together form a ring


WO 2005/051931

PCT/IB2004/003893

73


27 fused with the ring A containing 3-5 carbon atoms within the ring and having 1-3
28 heteroatoms selected from N, O or S;

29 reacting the compound of Formula III with hydroxylamine hydrochloride to give a
30 compound of Formula IV;




Formula IV

37 treating the compound of Formula IV with a compound of Formula V to give a
38 compound of Formula VI



Formula V



Formula VI

44 wherein

45 R₁ is selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl; cyano; nitro; amino;
46 substituted amino; hydroxyl; alkoxy; aryloxy; COR'; COOR'

47 (wherein R' can be hydrogen, alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl,
48 aryl, aralkyl, heterocyclyl, (heterocyclyl)alkyl, or (heteroaryl)alkyl);

49 aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl) alkyl; (heterocyclyl) alkyl; (CH₂)₁₋₄OR'

50 (wherein R' is as defined above, but also including hydroxy); C(=O)NR_xR_y,

51 (wherein R_x and R_y can be independently selected from hydrogen, alkyl, C₃₋₆

52 alkenyl, C₃₋₆ alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heteroaryl,

53 heterocyclyl, heteroarylalkyl, or heterocyclylalkyl); or

54 (CH₂)_m-C(=O)R₃

55 [wherein m is an integer in the range of 0-2 and R₃ can be optionally substituted

WO 2005/051931

PCT/IB2004/003893


74

56 R_p or R_q (wherein R_p can be a 4-12 membered (un)saturated monocyclic or
57 bicyclic ring containing 1-4 heteroatom(s) selected from N, O and S wherein the
58 ring can be attached to $(CH_2)_mC(=O)$ through N and R_q can be a 4-12 membered
59 (un)saturated monocyclic or bicyclic ring containing 0-4 heteroatom(s) selected
60 from the group consisting of N, O and S wherein the ring can be attached to
61 $(CH_2)_mC(=O)$ through C) and wherein the substituents of R_3 can be one or more
62 of: alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl, halogen, hydroxyl, alkoxy,
63 aryloxy, nitro, cyano, amino, substituted amino, hydroxyalkyl, oxo, acyl,
64 optionally substituted amino (wherein the substituents are selected from C_1-C_6
65 alkyl, aryl, aralkyl, or cycloalkyl), aryl, carboxyl, alkaryl, carbamoyl, alkyl ether,
66 $C(=O)NR_5R_6$ (wherein R_5 and R_6 are independently selected from hydrogen,
67 alkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, aryl, and aralkyl), optionally substituted
68 monocyclic or bicyclic 4-12 membered carbocyclic ring system (wherein the
69 optional substituent(s) is/are selected from alkyl, alkenyl, alkynyl, halogen,
70 hydroxyl, and alkoxy), heteroaryl, heterocyclyl, heteroarylalkyl, or
71 heterocyclylalkyl];

72 R_4 is selected from: hydrogen; alkyl; halogen; cyano; carboxy; or $C(=O)NR_xR_y$ wherein
73 R_x and R_y are the same as defined above;

74 and R_r represents $[(CH_2)_nCN, COOH, COOCH_3, CHO$ or pyridyl, wherein n is 0 to
75 2)];

76 reacting the compound of Formula VI with hydroxylamine hydrochloride (when R_r
77 is CN) to give a compound of Formula VII; and



Formula VII

84 reacting the compound of Formula VII with a compound of Formula $(R'CO)_2O$ to
85 give the compound of Formula VII(a) (wherein R' can be hydrogen, alkyl, alkenyl,


WO 2005/051931

PCT/IB2004/003893

75


86 alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heterocyclyl, (heterocyclyl)alkyl, or
87 (heteroaryl)alkyl).

1 14. A method for the preparation of compounds of Formula IX,



7 their pharmaceutically acceptable salts, pharmaceutically acceptable solvates,
8 enantiomers, diastereomers or N-oxides, the method comprising:

9 reacting a compound of Formula VI (when R_r is COOCH₃) with hydrazine hydrate
10 to give a compounds of Formula VIII



18 wherein

19 R₁ is selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl; cyano; nitro; amino;
20 substituted amino; hydroxyl; alkoxy; aryloxy; COR'; COOR'

21 (wherein R' can be hydrogen, alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl,
22 aryl, aralkyl, heterocyclyl, (heterocyclyl)alkyl, or (heteroaryl)alkyl);

23 aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl) alkyl; (heterocyclyl) alkyl; (CH₂)₁₋₄OR'

24 (wherein R' is as defined above, but also including hydroxy); C(=O)NR_xR_y

25 (wherein R_x and R_y can be independently selected from hydrogen, alkyl, C₃₋₆

26 alkenyl, C₃₋₆ alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heteroaryl,

27 heterocyclyl, heteroarylalkyl, or heterocyclylalkyl); or

WO 2005/051931

PCT/IB2004/003893

76

- 28 $(\text{CH}_2)_m\text{C}(=\text{O})\text{R}_3$
29 [wherein m is an integer in the range of 0-2 and R_3 can be optionally substituted
30 R_p or R_q (wherein R_p can be a 4-12 membered (un)saturated monocyclic or
31 bicyclic ring containing 1-4 heteroatom(s) selected from N, O and S wherein the
32 ring can be attached to $(\text{CH}_2)_m\text{C}(=\text{O})$ through N and R_q can be a 4-12 membered
33 (un)saturated monocyclic or bicyclic ring containing 0-4 heteroatom(s) selected
34 from the group consisting of N, O and S wherein the ring can be attached to
35 $(\text{CH}_2)_m\text{C}(=\text{O})$ through C) and wherein the substituents of R_3 can be one or more
36 of: alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl, halogen, hydroxyl, alkoxy,
37 aryloxy, nitro, cyano, amino, substituted amino, hydroxyalkyl, oxo, acyl,
38 optionally substituted amino (wherein the substituents are selected from $\text{C}_1\text{-C}_6$
39 alkyl, aryl, aralkyl, or cycloalkyl), aryl, carboxyl, alkaryl, carbamoyl, alkyl ether,
40 $\text{C}(=\text{O})\text{NR}_5\text{R}_6$ (wherein R_5 and R_6 are independently selected from hydrogen,
41 alkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, aryl, and aralkyl), optionally substituted
42 monocyclic or bicyclic 4-12 membered carbocyclic ring system (wherein the
43 optional substituent(s) is/are selected from alkyl, alkenyl, alkynyl, halogen,
44 hydroxyl, and alkoxy), heteroaryl, heterocyclyl, heteroarylalkyl, or
45 heterocyclylalkyl];
- 46 R_4 is selected from: hydrogen; alkyl; halogen; cyano; carboxy; or $\text{C}(=\text{O})\text{NR}_x\text{R}_y$ wherein
47 R_x and R_y are the same as defined above;
- 48 X_1 and X_2 are independently selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl;
49 acyl; aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl)alkyl; or (heterocyclyl)alkyl;
- 50 Y is selected from: an oxygen atom; a sulphur atom; or NR
51 (wherein R is selected from hydrogen, alkyl, alkenyl, alkynyl, un(saturated)
52 cycloalkyl, acyl, aryl, aralkyl, heteroaryl, heterocyclyl, (heteroaryl)alkyl, or
53 (heterocyclyl)alkyl);
- 54 Y_1 and Y_2 are independently selected from: hydrogen; alkyl; nitro; cyano; halogen; OR
55 wherein R is the same as defined earlier; SR wherein R is the same as defined earlier;
56 NHR wherein R is the same as defined earlier; COOR' ; or COR' wherein R' is the same
57 as defined above, or further, Y_1 and X_2 , X_1 and Y_2 , X_1 and X_2 may together form a ring

WO 2005/051931


PCT/IB2004/003893

77

58 fused with the ring A containing 3-5 carbon atoms within the ring and having 1-3
59 heteroatoms selected from N, O or S;

60 reacting the compound of Formula VIII with a compound of Formula $HC(OR_{11})_3$
61 to give a compound of Formula IX (wherein R_{11} represents alkyl from C_1 to C_3).


1 15. A method for the preparation of compounds of Formula X,



Formula X

7 their pharmaceutically acceptable salts, pharmaceutically acceptable solvates,
8 enantiomers, diastereomers or N-oxides, the method comprising:

9 reacting a compound of Formula VI (when R_r is CN)



Formula VI

15 wherein

16 R_1 is selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl; cyano; nitro; amino;
17 substituted amino; hydroxyl; alkoxy; aryloxy; COR' ; $COOR'$

18 (wherein R' can be hydrogen, alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl,
19 aryl, aralkyl, heterocyclyl, (heterocyclyl)alkyl, or (heteroaryl)alkyl);

20 aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl) alkyl; (heterocyclyl) alkyl; $(CH_2)_{1-4}OR'$

21 (wherein R' is as defined above, but also including hydroxy); $C(=O)NR_xR_y$

22 (wherein R_x and R_y can be independently selected from hydrogen, alkyl, C_{3-6}

23 alkenyl, C_{3-6} alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heteroaryl,

24 heterocyclyl, heteroarylalkyl, or heterocyclylalkyl); or

25 $(CH_2)_m-C(=O)R_3$

WO 2005/051931

PCT/IB2004/003893

78

26 [wherein m is an integer in the range of 0-2 and R₃ can be optionally substituted
27 R_p or R_q (wherein R_p can be a 4-12 membered (un)saturated monocyclic or
28 bicyclic ring containing 1-4 heteroatom(s) selected from N, O and S wherein the
29 ring can be attached to (CH₂)_mC(=O) through N and R_q can be a 4-12 membered
30 (un)saturated monocyclic or bicyclic ring containing 0-4 heteroatom(s) selected
31 from the group consisting of N, O and S wherein the ring can be attached to
32 (CH₂)_mC(=O) through C) and wherein the substituents of R₃ can be one or more
33 of: alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl, halogen, hydroxyl, alkoxy,
34 aryloxy, nitro, cyano, amino, substituted amino, hydroxyalkyl, oxo, acyl,
35 optionally substituted amino (wherein the substituents are selected from C₁-C₆
36 alkyl, aryl, aralkyl, or cycloalkyl), aryl, carboxyl, alkaryl, carbamoyl, alkyl ether,
37 C(=O)NR₅R₆ (wherein R₅ and R₆ are independently selected from hydrogen,
38 alkyl, C₃₋₆ alkenyl, C₃₋₆ alkynyl, aryl, and aralkyl), optionally substituted
39 monocyclic or bicyclic 4-12 membered carbocyclic ring system (wherein the
40 optional substituent(s) is/are selected from alkyl, alkenyl, alkynyl, halogen,
41 hydroxyl, and alkoxy), heteroaryl, heterocyclyl, heteroarylalkyl, or
42 heterocyclylalkyl];

43 R₄ is selected from: hydrogen; alkyl; halogen; cyano; carboxy; or C(=O)NR_xR_y wherein
44 R_x and R_y are the same as defined above;

45 X₁ and X₂ are independently selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl;
46 acyl; aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl)alkyl; or (heterocyclyl)alkyl;

47 Y₁ and Y₂ are independently selected from: hydrogen; alkyl; nitro; cyano; halogen; OR
48 wherein R is the same as defined earlier; SR wherein R is the same as defined earlier;
49 NHR wherein R is the same as defined earlier; COOR'; or COR' wherein R' is the same
50 as defined above, or further, Y₁ and X₂, X₁ and Y₂, X₁ and X₂ may together form a ring
51 fused with the ring A containing 3-5 carbon atoms within the ring and having 1-3
52 heteroatoms selected from N, O or S;


53 with sodium azide to give the compound of Formula X.

WO 2005/051931

PCT/IB2004/003893


79

16. A method for the preparation of compounds of Formula XI,



their pharmaceutically acceptable salts, pharmaceutically acceptable solvates, enantiomers, diastereomers or N-oxides, the method comprising:

reacting a compound of Formula VII



wherein

R₁ is selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl; cyano; nitro; amino; substituted amino; hydroxyl; alkoxy; aryloxy; COR'; COOR'

(wherein R' can be hydrogen, alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heterocyclyl, (heterocyclyl)alkyl, or (heteroaryl)alkyl);

aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl) alkyl; (heterocyclyl) alkyl; (CH₂)₁₋₄OR'

(wherein R' is as defined above, but also including hydroxy); C(=O)NR_xR_y

(wherein R_x and R_y can be independently selected from hydrogen, alkyl, C₃₋₆

alkenyl, C₃₋₆ alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heteroaryl,

heterocyclyl, heteroarylalkyl, or heterocyclylalkyl); or

(CH₂)_m-C(=O)R₃

[wherein m is an integer in the range of 0-2 and R₃ can be optionally substituted

R_p or R_q (wherein R_p can be a 4-12 membered (un)saturated monocyclic or

bicyclic ring containing 1-4 heteroatom(s) selected from N, O and S wherein the

WO 2005/051931

PCT/IB2004/003893

80

28 ring can be attached to $(CH_2)_mC(=O)$ through N and R_q can be a 4-12 membered
 29 (un)saturated monocyclic or bicyclic ring containing 0-4 heteroatom(s) selected
 30 from the group consisting of N, O and S wherein the ring can be attached to
 31 $(CH_2)_mC(=O)$ through C) and wherein the substituents of R_3 can be one or more
 32 of: alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl, halogen, hydroxyl, alkoxy,
 33 aryloxy, nitro, cyano, amino, substituted amino, hydroxyalkyl, oxo, acyl,
 34 optionally substituted amino (wherein the substituents are selected from C_1-C_6
 35 alkyl, aryl, aralkyl, or cycloalkyl), aryl, carboxyl, alkaryl, carbamoyl, alkyl ether,
 36 $C(=O)NR_5R_6$ (wherein R_5 and R_6 are independently selected from hydrogen,
 37 alkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, aryl, and aralkyl), optionally substituted
 38 monocyclic or bicyclic 4-12 membered carbocyclic ring system (wherein the
 39 optional substituent(s) is/are selected from alkyl, alkenyl, alkynyl, halogen,
 40 hydroxyl, and alkoxy), heteroaryl, heterocyclyl, heteroarylalkyl, or
 41 heterocyclylalkyl];


42 R_4 is selected from: hydrogen; alkyl; halogen; cyano; carboxy; or $C(=O)NR_xR_y$ wherein
 43 R_x and R_y are the same as defined above;

44 X_1 and X_2 are independently selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl;
 45 acyl; aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl)alkyl; or (heterocyclyl)alkyl;

46 Y_1 and Y_2 are independently selected from: hydrogen; alkyl; nitro; cyano; halogen; OR
 47 wherein R is the same as defined earlier; SR wherein R is the same as defined earlier;
 48 NHR wherein R is the same as defined earlier; $COOR'$; or COR' wherein R' is the same
 49 as defined above, or further, Y_1 and X_2 , X_1 and Y_2 , X_1 and X_2 may together form a ring
 50 fused with the ring A containing 3-5 carbon atoms within the ring and having 1-3
 51 heteroatoms selected from N, O or S;


52 with methyl chloroformate to give the compound of Formula XI.

17. A method for the preparation of compounds of Formula XII,



their pharmaceutically acceptable salts, pharmaceutically acceptable solvates, enantiomers, diastereomers or N-oxides, the method comprising:

reacting compounds of Formula VII



wherein

R₁ is selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl; cyano; nitro; amino; substituted amino; hydroxyl; alkoxy; aryloxy; COR'; COOR' (wherein R' can be hydrogen, alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heterocyclyl, (heterocyclyl)alkyl, or (heteroaryl)alkyl); aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl) alkyl; (heterocyclyl) alkyl; (CH₂)₁₋₄OR' (wherein R' is as defined above, but also including hydroxy); C(=O)NR_xR_y (wherein R_x and R_y can be independently selected from hydrogen, alkyl, C₃₋₆ alkenyl, C₃₋₆ alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heteroaryl, heterocyclyl, heteroarylalkyl, or heterocyclylalkyl); or (CH₂)_m-C(=O)R₃ [wherein m is an integer in the range of 0-2 and R₃ can be optionally substituted R_p or R_q (wherein R_p can be a 4-12 membered (un)saturated monocyclic or bicyclic ring containing 1-4 heteroatom(s) selected from N, O and S wherein the ring can be attached to (CH₂)_mC(=O) through N and R_q can be a 4-12 membered

WO 2005/051931

PCT/IB2004/003893

82

30 (un)saturated monocyclic or bicyclic ring containing 0-4 heteroatom(s) selected
31 from the group consisting of N, O and S wherein the ring can be attached to
32 $(\text{CH}_2)_m\text{C}(=\text{O})$ through C) and wherein the substituents of R_3 can be one or more
33 of: alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl, halogen, hydroxyl, alkoxy,
34 aryloxy, nitro, cyano, amino, substituted amino, hydroxyalkyl, oxo, acyl,
35 optionally substituted amino (wherein the substituents are selected from $\text{C}_1\text{-C}_6$
36 alkyl, aryl, aralkyl, or cycloalkyl), aryl, carboxyl, alkaryl, carbamoyl, alkyl ether,
37 $\text{C}(=\text{O})\text{NR}_5\text{R}_6$ (wherein R_5 and R_6 are independently selected from hydrogen,
38 alkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, aryl, and aralkyl), optionally substituted
39 monocyclic or bicyclic 4-12 membered carbocyclic ring system (wherein the
40 optional substituent(s) is/are selected from alkyl, alkenyl, alkynyl, halogen,
41 hydroxyl, and alkoxy), heteroaryl, heterocyclyl, heteroarylalkyl, or
42 heterocyclylalkyl];

43 R_4 is selected from: hydrogen; alkyl; halogen; cyano; carboxy; or $\text{C}(=\text{O})\text{NR}_x\text{R}_y$ wherein
44 R_x and R_y are the same as defined above;

45 X_1 and X_2 are independently selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl;
46 acyl; aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl)alkyl; or (heterocyclyl)alkyl;

47 Y_1 and Y_2 are independently selected from: hydrogen; alkyl; nitro; cyano; halogen; OR
48 wherein R is the same as defined earlier; SR wherein R is the same as defined earlier;
49 NHR wherein R is the same as defined earlier; COOR' ; or COR' wherein R' is the same
50 as defined above, or further, Y_1 and X_2 , X_1 and Y_2 , X_1 and X_2 may together form a ring
51 fused with the ring A containing 3-5 carbon atoms within the ring and having 1-3
52 heteroatoms selected from N, O or S;


53 with thiocarbonyl diimidazole and 1,8-diazabicyclo[5.4.0]undec-7-one to give the
54 compound of Formula XII.

WO 2005/051931

PCT/IB2004/003893


83

18. A method for the preparation of compounds of Formula XIII,



their pharmaceutically acceptable salts, pharmaceutically acceptable solvates, enantiomers, diastereomers or N-oxides, the method comprising:

treating a compounds of Formula XII,



wherein

R₁ is selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl; cyano; nitro; amino; substituted amino; hydroxyl; alkoxy; aryloxy; COR'; COOR'

(wherein R' can be hydrogen, alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heterocyclyl, (heterocyclyl)alkyl, or (heteroaryl)alkyl);

aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl) alkyl; (heterocyclyl) alkyl; (CH₂)₁₋₄OR'

(wherein R' is as defined above, but also including hydroxy); C(=O)NR_xR_y

(wherein R_x and R_y can be independently selected from hydrogen, alkyl, C₃₋₆ alkenyl, C₃₋₆ alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heteroaryl, heterocyclyl, heteroarylalkyl, or heterocyclylalkyl); or

(CH₂)_m-C(=O)R₃

[wherein m is an integer in the range of 0-2 and R₃ can be optionally substituted

R_p or R_q (wherein R_p can be a 4-12 membered (un)saturated monocyclic or

bicyclic ring containing 1-4 heteroatom(s) selected from N, O and S wherein the

ring can be attached to (CH₂)_mC(=O) through N and R_q can be a 4-12 membered

WO 2005/051931

PCT/IB2004/003893

84

31 (un)saturated monocyclic or bicyclic ring containing 0-4 heteroatom(s) selected
32 from the group consisting of N, O and S wherein the ring can be attached to
33 $(CH_2)_mC(=O)$ through C) and wherein the substituents of R_3 can be one or more
34 of: alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl, halogen, hydroxyl, alkoxy,
35 aryloxy, nitro, cyano, amino, substituted amino, hydroxyalkyl, oxo, acyl,
36 optionally substituted amino (wherein the substituents are selected from C_1-C_6
37 alkyl, aryl, aralkyl, or cycloalkyl), aryl, carboxyl, alkaryl, carbamoyl, alkyl ether,
38 $C(=O)NR_5R_6$ (wherein R_5 and R_6 are independently selected from hydrogen,
39 alkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, aryl, and aralkyl), optionally substituted
40 monocyclic or bicyclic 4-12 membered carbocyclic ring system (wherein the
41 optional substituent(s) is/are selected from alkyl, alkenyl, alkynyl, halogen,
42 hydroxyl, and alkoxy), heteroaryl, heterocyclyl, heteroarylalkyl, or
43 heterocyclylalkyl];

44 R_4 is selected from: hydrogen; alkyl; halogen; cyano; carboxy; or $C(=O)NR_xR_y$ wherein
45 R_x and R_y are the same as defined above;

46 X_1 and X_2 are independently selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl;
47 acyl; aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl)alkyl; or (heterocyclyl)alkyl;

48 Y_1 and Y_2 are independently selected from: hydrogen; alkyl; nitro; cyano; halogen; OR
49 wherein R is the same as defined earlier; SR wherein R is the same as defined earlier;
50 NHR wherein R is the same as defined earlier; $COOR'$; or COR' wherein R' is the same
51 as defined above, or further, Y_1 and X_2 , X_1 and Y_2 , X_1 and X_2 may together form a ring
52 fused with the ring A containing 3-5 carbon atoms within the ring and having 1-3
53 heteroatoms selected from N, O or S;


54 with a compound of Formula $R_{11}Z$ (wherein Z is halogen) to gives the compound
55 of Formula XIII (wherein R_{11} is alkyl).

WO 2005/051931

PCT/IB2004/003893


85

19. A method for the preparation of compounds of Formula XIV,



their pharmaceutically acceptable salts, pharmaceutically acceptable solvates,
enantiomers, diastereomers or N-oxides, the method comprising:

reacting a compound of Formula VII



wherein

R_1 is selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl; cyano; nitro; amino;
substituted amino; hydroxyl; alkoxy; aryloxy; COR' ; $COOR'$
(wherein R' can be hydrogen, alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl,
aryl, aralkyl, heterocyclyl, (heterocyclyl)alkyl, or (heteroaryl)alkyl);
aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl) alkyl; (heterocyclyl) alkyl; $(CH_2)_{1-4}OR'$
(wherein R' is as defined above, but also including hydroxy); $C(=O)NR_xR_y$
(wherein R_x and R_y can be independently selected from hydrogen, alkyl, C_{3-6}
alkenyl, C_{3-6} alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heteroaryl,
heterocyclyl, heteroarylalkyl, or heterocyclylalkyl); or
 $(CH_2)_m-C(=O)R_3$
[wherein m is an integer in the range of 0-2 and R_3 can be optionally substituted

WO 2005/051931

PCT/IB2004/003893

86

22 R_p or R_q (wherein R_p can be a 4-12 membered (un)saturated monocyclic or
23 bicyclic ring containing 1-4 heteroatom(s) selected from N, O and S wherein the
24 ring can be attached to $(CH_2)_mC(=O)$ through N and R_q can be a 4-12 membered
25 (un)saturated monocyclic or bicyclic ring containing 0-4 heteroatom(s) selected
26 from the group consisting of N, O and S wherein the ring can be attached to
27 $(CH_2)_mC(=O)$ through C) and wherein the substituents of R_3 can be one or more
28 of: alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl, halogen, hydroxyl, alkoxy,
29 aryloxy, nitro, cyano, amino, substituted amino, hydroxyalkyl, oxo, acyl,
30 optionally substituted amino (wherein the substituents are selected from C_1 - C_6
31 alkyl, aryl, aralkyl, or cycloalkyl), aryl, carboxyl, alkaryl, carbamoyl, alkyl ether,
32 $C(=O)NR_5R_6$ (wherein R_5 and R_6 are independently selected from hydrogen,
33 alkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, aryl, and aralkyl), optionally substituted
34 monocyclic or bicyclic 4-12 membered carbocyclic ring system (wherein the
35 optional substituent(s) is/are selected from alkyl, alkenyl, alkynyl, halogen,
36 hydroxyl, and alkoxy), heteroaryl, heterocyclyl, heteroarylalkyl, or
37 heterocyclylalkyl];

38 R_4 is selected from: hydrogen; alkyl; halogen; cyano; carboxy; or $C(=O)NR_xR_y$ wherein
39 R_x and R_y are the same as defined above;
40 X_1 and X_2 are independently selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl;
41 acyl; aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl)alkyl; or (heterocyclyl)alkyl;
42 Y_1 and Y_2 are independently selected from: hydrogen; alkyl; nitro; cyano; halogen; OR
43 wherein R is the same as defined earlier; SR wherein R is the same as defined earlier;
44 NHR wherein R is the same as defined earlier; $COOR'$; or COR' wherein R' is the same
45 as defined above, or further, Y_1 and X_2 , X_1 and Y_2 , X_1 and X_2 may together form a ring
46 fused with the ring A containing 3-5 carbon atoms within the ring and having 1-3
47 heteroatoms selected from N, O or S;


48 with thiocarbonyl diimidazole and boron trifluoride etherate to give the compound
49 of Formula XIV.

WO 2005/051931


PCT/IB2004/003893

87

20. A method for the preparation of compounds of Formula XV,



their pharmaceutically acceptable salts, pharmaceutically acceptable solvates,
enantiomers, diastereomers or N-oxides, the method comprising:
reacting compounds of Formula VII



wherein

R_1 is selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl; cyano; nitro; amino;
substituted amino; hydroxyl; alkoxy; aryloxy; COR'; COOR'

(wherein R' can be hydrogen, alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl,
aryl, aralkyl, heterocyclyl, (heterocyclyl)alkyl, or (heteroaryl)alkyl);

aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl) alkyl; (heterocyclyl) alkyl; $(CH_2)_{1-4}OR'$
(wherein R' is as defined above, but also including hydroxy); $C(=O)NR_xR_y$
(wherein R_x and R_y can be independently selected from hydrogen, alkyl, C_{3-6}
alkenyl, C_{3-6} alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heteroaryl,
heterocyclyl, heteroarylalkyl, or heterocyclylalkyl); or
 $(CH_2)_m-C(=O)R_3$
[wherein m is an integer in the range of 0-2 and R_3 can be optionally substituted
 R_p or R_q (wherein R_p can be a 4-12 membered (un)saturated monocyclic or
bicyclic ring containing 1-4 heteroatom(s) selected from N, O and S wherein the

WO 2005/051931

PCT/IB2004/003893

88

26 ring can be attached to $(CH_2)_mC(=O)$ through N and R_q can be a 4-12 membered
27 (un)saturated monocyclic or bicyclic ring containing 0-4 heteroatom(s) selected
28 from the group consisting of N, O and S wherein the ring can be attached to
29 $(CH_2)_mC(=O)$ through C) and wherein the substituents of R_3 can be one or more
30 of: alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl, halogen, hydroxyl, alkoxy,
31 aryloxy, nitro, cyano, amino, substituted amino, hydroxyalkyl, oxo, acyl,
32 optionally substituted amino (wherein the substituents are selected from C_1-C_6
33 alkyl, aryl, aralkyl, or cycloalkyl), aryl, carboxyl, alkaryl, carbamoyl, alkyl ether,
34 $C(=O)NR_5R_6$ (wherein R_5 and R_6 are independently selected from hydrogen,
35 alkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, aryl, and aralkyl), optionally substituted
36 monocyclic or bicyclic 4-12 membered carbocyclic ring system (wherein the
37 optional substituent(s) is/are selected from alkyl, alkenyl, alkynyl, halogen,
38 hydroxyl, and alkoxy), heteroaryl, heterocyclyl, heteroarylalkyl, or
39 heterocyclalkyl];

40 R_4 is selected from: hydrogen; alkyl; halogen; cyano; carboxy; or $C(=O)NR_xR_y$ wherein
41 R_x and R_y are the same as defined above;

42 X_1 and X_2 are independently selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl;
43 acyl; aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl)alkyl; or (heterocyclyl)alkyl;

44 Y_1 and Y_2 are independently selected from: hydrogen; alkyl; nitro; cyano; halogen; OR
45 wherein R is the same as defined earlier; SR wherein R is the same as defined earlier;
46 NHR wherein R is the same as defined earlier; COOR'; or COR' wherein R' is the same
47 as defined above, or further, Y_1 and X_2 , X_1 and Y_2 , X_1 and X_2 may together form a ring
48 fused with the ring A containing 3-5 carbon atoms within the ring and having 1-3
49 heteroatoms selected from N, O or S;


50 with compounds of Formula (a) $R_{12}COOH$; (b) $R_{12}COCl$ or (c) $R_{12}COOC_2H_5$ to
51 give the compound of Formula XV (wherein R_{12} is alkyl, cycloalkyl, aryl,
52 heteroaryl or heterocyclyl).

WO 2005/051931

PCT/IB2004/003893

89

21. A method for the preparation of compounds of Formula XX,



their pharmaceutically acceptable salts, pharmaceutically acceptable solvates,
enantiomers, diastereomers or N-oxides,
wherein

R₁ is selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl; cyano; nitro; amino;
substituted amino; hydroxyl; alkoxy; aryloxy; COR'; COOR'

(wherein R' can be hydrogen, alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl,
aryl, aralkyl, heterocyclyl, (heterocyclyl)alkyl, or (heteroaryl)alkyl);

aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl) alkyl; (heterocyclyl) alkyl; (CH₂)₁₋₄OR'
(wherein R' is as defined above, but also including hydroxy); C(=O)NR_xR_y

(wherein R_x and R_y can be independently selected from hydrogen, alkyl, C₃₋₆
alkenyl, C₃₋₆ alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heteroaryl,
heterocyclyl, heteroarylalkyl, or heterocyclylalkyl); or

(CH₂)_m-C(=O)R₃

[wherein m is an integer in the range of 0-2 and R₃ can be optionally substituted
R_p or R_q (wherein R_p can be a 4-12 membered (un)saturated monocyclic or
bicyclic ring containing 1-4 heteroatom(s) selected from N, O and S wherein the
ring can be attached to (CH₂)_mC(=O) through N and R_q can be a 4-12 membered
(un)saturated monocyclic or bicyclic ring containing 0-4 heteroatom(s) selected
from the group consisting of N, O and S wherein the ring can be attached to
(CH₂)_mC(=O) through C) and wherein the substituents of R₃ can be one or more
of: alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl, halogen, hydroxyl, alkoxy,
aryloxy, nitro, cyano, amino, substituted amino, hydroxyalkyl, oxo, acyl,
optionally substituted amino (wherein the substituents are selected from C₁-C₆
alkyl, aryl, aralkyl, or cycloalkyl), aryl, carboxyl, alkaryl, carbamoyl, alkyl ether,
C(=O)NR₅R₆ (wherein R₅ and R₆ are independently selected from hydrogen,


WO 2005/051931

PCT/IB2004/003893


90

32 alkyl, C₃₋₆ alkenyl, C₃₋₆ alkynyl, aryl, and aralkyl), optionally substituted
 33 monocyclic or bicyclic 4-12 membered carbocyclic ring system (wherein the
 34 optional substituent(s) is/are selected from alkyl, alkenyl, alkynyl, halogen,
 35 hydroxyl, and alkoxy), heteroaryl, heterocyclyl, heteroarylalkyl, or
 36 heterocyclylalkyl];
 37 R₄ is selected from: hydrogen; alkyl; halogen; cyano; carboxy; or C(=O)NR_xR_y wherein
 38 R_x and R_y are the same as defined above;
 39 X₁ and X₂ are independently selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl;
 40 acyl; aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl)alkyl; or (heterocyclyl)alkyl;
 41 Y₁ and Y₂ are independently selected from: hydrogen; alkyl; nitro; cyano; halogen; OR
 42 wherein R is the same as defined earlier; SR wherein R is the same as defined earlier;
 43 NHR wherein R is the same as defined earlier; COOR'; or COR' wherein R' is the same
 44 as defined above, or further, Y₁ and X₂, X₁ and Y₂, X₁ and X₂ may together form a ring
 45 fused with the ring A containing 3-5 carbon atoms within the ring and having 1-3
 46 heteroatoms selected from N, O or S; and
 47 R₁₂ is alkyl, cycloalkyl, aryl, heteroaryl or heterocyclyl;

48 the method comprising:
 49 reacting a compound of Formula IV with a compound of Formula XVI



to give a compound of Formula XVII;




WO 2005/051931


PCT/IB2004/003893

91

treating the compound of Formula XVII with potassium phthalamide to give a compound of Formula XVIII;




treating the compound of Formula XVIII with a hydrazine hydrate to give a compound of Formula XIX; and



treating the compound of Formula XIX with a compound of Formula $R_{12}COCl$ or $R_{12}COOH$ to give the compound of Formula XX.

22. A method for the preparation of compounds of Formula XXIII,



their pharmaceutically acceptable salts, pharmaceutically acceptable solvates, enantiomers, diastereomers or N-oxides, wherein

R_1 is selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl; cyano; nitro; amino; substituted amino; hydroxyl; alkoxy; aryloxy; COR' ; $COOR'$

(wherein R' can be hydrogen, alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl,

WO 2005/051931


PCT/IB2004/003893

92


14 aryl, aralkyl, heterocyclyl, (heterocyclyl)alkyl, or (heteroaryl)alkyl);
 15 aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl) alkyl; (heterocyclyl) alkyl; (CH₂)₁₋₄OR'
 16 (wherein R' is as defined above, but also including hydroxy); C(=O)NR_xR_y
 17 (wherein R_x and R_y can be independently selected from hydrogen, alkyl, C₃₋₆
 18 alkenyl, C₃₋₆ alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heteroaryl,
 19 heterocyclyl, heteroarylalkyl, or heterocyclylalkyl); or
 20 (CH₂)_m-C(=O)R₃
 21 [wherein m is an integer in the range of 0-2 and R₃ can be optionally substituted
 22 R_p or R_q (wherein R_p can be a 4-12 membered (un)saturated monocyclic or
 23 bicyclic ring containing 1-4 heteroatom(s) selected from N, O and S wherein the
 24 ring can be attached to (CH₂)_mC(=O) through N and R_q can be a 4-12 membered
 25 (un)saturated monocyclic or bicyclic ring containing 0-4 heteroatom(s) selected
 26 from the group consisting of N, O and S wherein the ring can be attached to
 27 (CH₂)_mC(=O) through C) and wherein the substituents of R₃ can be one or more
 28 of: alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl, halogen, hydroxyl, alkoxy,
 29 aryloxy, nitro, cyano, amino, substituted amino, hydroxyalkyl, oxo, acyl,
 30 optionally substituted amino (wherein the substituents are selected from C₁-C₆
 31 alkyl, aryl, aralkyl, or cycloalkyl), aryl, carboxyl, alkaryl, carbamoyl, alkyl ether,
 32 C(=O)NR₅R₆ (wherein R₅ and R₆ are independently selected from hydrogen,
 33 alkyl, C₃₋₆ alkenyl, C₃₋₆ alkynyl, aryl, and aralkyl), optionally substituted
 34 monocyclic or bicyclic 4-12 membered carbocyclic ring system (wherein the
 35 optional substituent(s) is/are selected from alkyl, alkenyl, alkynyl, halogen,
 36 hydroxyl, and alkoxy), heteroaryl, heterocyclyl, heteroarylalkyl, or
 37 heterocyclylalkyl];
 38 R₄ is selected from: hydrogen; alkyl; halogen; cyano; carboxy; or C(=O)NR_xR_y wherein
 39 R_x and R_y are the same as defined above;
 40 X₁ and X₂ are independently selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl;
 41 acyl; aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl)alkyl; or (heterocyclyl)alkyl;
 42 Y₁ and Y₂ are independently selected from: hydrogen; alkyl; nitro; cyano; halogen; OR
 43 wherein R is the same as defined earlier; SR wherein R is the same as defined earlier;
 44 NHR wherein R is the same as defined earlier; COOR'; or COR' wherein R' is the same

45 as defined above, or further, Y₁ and X₂, X₁ and Y₂, X₁ and X₂ may together form a ring
 46 fused with the ring A containing 3-5 carbon atoms within the ring and having 1-3
 47 heteroatoms selected from N, O or S; and
 48 R₁₃ is alkyl, aryl or heteroaryl;

49 the method comprising
 50 reacting compounds of Formula XXI with hydroxylamine hydrochloride to give
 51 compounds of Formula XXII ,




55 which on reaction with compounds of Formula VI (when R_r is COOH),



62 gives compounds of Formula XXIII.

1 23. A method for the preparation of compounds of Formula XXIV,



7 their pharmaceutically acceptable salts, pharmaceutically acceptable solvates,
 8 enantiomers, diastereomers or N-oxides,
 9 wherein
 10 R₁ is selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl; cyano; nitro; amino;
 11 substituted amino; hydroxyl; alkoxy; aryloxy; COR'; COOR'
 12 (wherein R' can be hydrogen, alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl,

WO 2005/051931

PCT/IB2004/003893

94

13 aryl, aralkyl, heterocyclyl, (heterocyclyl)alkyl, or (heteroaryl)alkyl);
 14 aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl) alkyl; (heterocyclyl) alkyl; (CH₂)₁₋₄OR'
 15 (wherein R' is as defined above, but also including hydroxy); C(=O)NR_xR_y
 16 (wherein R_x and R_y can be independently selected from hydrogen, alkyl, C₃₋₆
 17 alkenyl, C₃₋₆ alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heteroaryl,
 18 heterocyclyl, heteroarylalkyl, or heterocyclylalkyl); or
 19 (CH₂)_m-C(=O)R₃
 20 [wherein m is an integer in the range of 0-2 and R₃ can be optionally substituted
 21 R_p or R_q (wherein R_p can be a 4-12 membered (un)saturated monocyclic or
 22 bicyclic ring containing 1-4 heteroatom(s) selected from N, O and S wherein the
 23 ring can be attached to (CH₂)_mC(=O) through N and R_q can be a 4-12 membered
 24 (un)saturated monocyclic or bicyclic ring containing 0-4 heteroatom(s) selected
 25 from the group consisting of N, O and S wherein the ring can be attached to
 26 (CH₂)_mC(=O) through C) and wherein the substituents of R₃ can be one or more
 27 of: alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl, halogen, hydroxyl, alkoxy,
 28 aryloxy, nitro, cyano, amino, substituted amino, hydroxyalkyl, oxo, acyl,
 29 optionally substituted amino (wherein the substituents are selected from C₁-C₆
 30 alkyl, aryl, aralkyl, or cycloalkyl), aryl, carboxyl, alkaryl, carbamoyl, alkyl ether,
 31 C(=O)NR₅R₆ (wherein R₅ and R₆ are independently selected from hydrogen,
 32 alkyl, C₃₋₆ alkenyl, C₃₋₆ alkynyl, aryl, and aralkyl), optionally substituted
 33 monocyclic or bicyclic 4-12 membered carbocyclic ring system (wherein the
 34 optional substituent(s) is/are selected from alkyl, alkenyl, alkynyl, halogen,
 35 hydroxyl, and alkoxy), heteroaryl, heterocyclyl, heteroarylalkyl, or
 36 heterocyclylalkyl];
 37 R₄ is selected from: hydrogen; alkyl; halogen; cyano; carboxy; or C(=O)NR_xR_y wherein
 38 R_x and R_y are the same as defined above;
 39 X₁ and X₂ are independently selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl;
 40 acyl; aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl)alkyl; or (heterocyclyl)alkyl;
 41 Y₁ and Y₂ are independently selected from: hydrogen; alkyl; nitro; cyano; halogen; OR
 42 wherein R is the same as defined earlier; SR wherein R is the same as defined earlier;
 43 NHR wherein R is the same as defined earlier; COOR'; or COR' wherein R' is the same

WO 2005/051931


PCT/IB2004/003893

95

as defined above, or further, Y_1 and X_2 , X_1 and Y_2 , X_1 and X_2 may together form a ring fused with the ring A containing 3-5 carbon atoms within the ring and having 1-3 heteroatoms selected from N, O or S;

the method comprising:


reacting a compound of Formula VI (when Rr is CN)



Formula VI

with $NH_2CH_2CH_2SH \cdot HCl$ to give the compounds of Formula XXIV.

24. A method for the preparation of compounds of Formula XXV,



Formula XXV

their pharmaceutically acceptable salts, pharmaceutically acceptable solvates, enantiomers, diastereomers or N-oxides, wherein

R_1 is selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl; cyano; nitro; amino; substituted amino; hydroxyl; alkoxy; aryloxy; COR' ; $COOR'$

(wherein R' can be hydrogen, alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heterocyclyl, (heterocyclyl)alkyl, or (heteroaryl)alkyl);

aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl) alkyl; (heterocyclyl) alkyl; $(CH_2)_{1-4}OR'$

(wherein R' is as defined above, but also including hydroxy); $C(=O)NR_xR_y$

(wherein R_x and R_y can be independently selected from hydrogen, alkyl, C_{3-6}

alkenyl, C_{3-6} alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heteroaryl,

heterocyclyl, heteroarylalkyl, or heterocyclylalkyl); or

$(CH_2)_m-C(=O)R_3$

[wherein m is an integer in the range of 0-2 and R_3 can be optionally substituted

WO 2005/051931

PCT/IB2004/003893

96

75 R_p or R_q (wherein R_p can be a 4-12 membered (un)saturated monocyclic or
76 bicyclic ring containing 1-4 heteroatom(s) selected from N, O and S wherein the
77 ring can be attached to $(CH_2)_mC(=O)$ through N and R_q can be a 4-12 membered
78 (un)saturated monocyclic or bicyclic ring containing 0-4 heteroatom(s) selected
79 from the group consisting of N, O and S wherein the ring can be attached to
80 $(CH_2)_mC(=O)$ through C) and wherein the substituents of R_3 can be one or more
81 of: alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl, halogen, hydroxyl, alkoxy,
82 aryloxy, nitro, cyano, amino, substituted amino, hydroxyalkyl, oxo, acyl,
83 optionally substituted amino (wherein the substituents are selected from C_1-C_6
84 alkyl, aryl, aralkyl, or cycloalkyl), aryl, carboxyl, alkaryl, carbamoyl, alkyl ether,
85 $C(=O)NR_5R_6$ (wherein R_5 and R_6 are independently selected from hydrogen,
86 alkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, aryl, and aralkyl), optionally substituted
87 monocyclic or bicyclic 4-12 membered carbocyclic ring system (wherein the
88 optional substituent(s) is/are selected from alkyl, alkenyl, alkynyl, halogen,
89 hydroxyl, and alkoxy), heteroaryl, heterocyclyl, heteroarylalkyl, or
90 heterocyclylalkyl];


91 R_4 is selected from: hydrogen; alkyl; halogen; cyano; carboxy; or $C(=O)NR_xR_y$ wherein
92 R_x and R_y are the same as defined above;

93 X_1 and X_2 are independently selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl;
94 acyl; aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl)alkyl; or (heterocyclyl)alkyl;

95 Y_1 and Y_2 are independently selected from: hydrogen; alkyl; nitro; cyano; halogen; OR
96 wherein R is the same as defined earlier; SR wherein R is the same as defined earlier;
97 NHR wherein R is the same as defined earlier; COOR'; or COR' wherein R' is the same
98 as defined above, or further, Y_1 and X_2 , X_1 and Y_2 , X_1 and X_2 may together form a ring
99 fused with the ring A containing 3-5 carbon atoms within the ring and having 1-3
100 heteroatoms selected from N, O or S;

101 the method comprising:


102 reacting a Formula VI



Formula VI

103
104 (wherein Rr is COOH) with NH₂NHCSNHR₁₄ (wherein R₁₄ represents hydrogen,
105 alkyl or cycloalkyl) to give the compound of Formula XXV.

1 25. A method for the preparation of compounds of Formula XXVII,



Formula XXVII

8 their pharmaceutically acceptable salts, pharmaceutically acceptable solvates,
9 enantiomers, diastereomers or N-oxides,
10 wherein
11 R₁ is selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl; cyano; nitro; amino;
12 substituted amino; hydroxyl; alkoxy; aryloxy, COR'; COOR'
13 (wherein R' can be hydrogen, alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl,
14 aryl, aralkyl, heterocyclyl, (heterocyclyl)alkyl, or (heteroaryl)alkyl);
15 aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl) alkyl; (heterocyclyl) alkyl; (CH₂)₁₋₄OR'
16 (wherein R' is as defined above, but also including hydroxy); C(=O)NR_xR_y
17 (wherein R_x and R_y can be independently selected from hydrogen, alkyl, C₃₋₆
18 alkenyl, C₃₋₆ alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heteroaryl,
19 heterocyclyl, heteroarylalkyl, or heterocyclylalkyl); or
20 (CH₂)_m-C(=O)R₃
21 [wherein m is an integer in the range of 0-2 and R₃ can be optionally substituted
22 R_p or R_q (wherein R_p can be a 4-12 membered (un)saturated monocyclic or

WO 2005/051931

PCT/IB2004/003893

98

23 bicyclic ring containing 1-4 heteroatom(s) selected from N, O and S wherein the
 24 ring can be attached to $(CH_2)_mC(=O)$ through N and R_q can be a 4-12 membered
 25 (un)saturated monocyclic or bicyclic ring containing 0-4 heteroatom(s) selected
 26 from the group consisting of N, O and S wherein the ring can be attached to
 27 $(CH_2)_mC(=O)$ through C) and wherein the substituents of R_3 can be one or more
 28 of: alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl, halogen, hydroxyl, alkoxy,
 29 aryloxy, nitro, cyano, amino, substituted amino, hydroxyalkyl, oxo, acyl,
 30 optionally substituted amino (wherein the substituents are selected from C_1-C_6
 31 alkyl, aryl, aralkyl, or cycloalkyl), aryl, carboxyl, alkaryl, carbamoyl, alkyl ether,
 32 $C(=O)NR_5R_6$ (wherein R_5 and R_6 are independently selected from hydrogen,
 33 alkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, aryl, and aralkyl), optionally substituted
 34 monocyclic or bicyclic 4-12 membered carbocyclic ring system (wherein the
 35 optional substituent(s) is/are selected from alkyl, alkenyl, alkynyl, halogen,
 36 hydroxyl, and alkoxy), heteroaryl, heterocyclyl, heteroarylalkyl, or
 37 heterocyclalkyl];

38 R_4 is selected from: hydrogen; alkyl; halogen; cyano; carboxy; or $C(=O)NR_xR_y$ wherein
 39 R_x and R_y are the same as defined above;


40 X_1 and X_2 are independently selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl;
 41 acyl; aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl)alkyl; or (heterocyclyl)alkyl;

42 Y_1 and Y_2 are independently selected from: hydrogen; alkyl; nitro; cyano; halogen; OR
 43 wherein R is the same as defined earlier; SR wherein R is the same as defined earlier;

44 NHR wherein R is the same as defined earlier; COOR'; or COR' wherein R' is the same
 45 as defined above, or further, Y_1 and X_2 , X_1 and Y_2 , X_1 and X_2 may together form a ring
 46 fused with the ring A containing 3-5 carbon atoms within the ring and having 1-3
 47 heteroatoms selected from N, O or S;


48 the method comprising:

reacting a compound of Formula VI



Formula VI


(wherein Rr is CHO) with hydroxylamine hydrochloride to give a compound of Formula XXVI; and



Formula XXVI

reacting the compound of Formula XXVI with methacrylonitrile to give the compound of Formula XXVII.

26. A method for the preparation of compounds of Formula XXIX,



Formula XXIX

their pharmaceutically acceptable salts, pharmaceutically acceptable solvates, enantiomers, diastereomers or N-oxides, wherein

R₁ is selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl; cyano; nitro; amino; substituted amino; hydroxyl; alkoxy; aryloxy; COR'; COOR' (wherein R' can be hydrogen, alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heterocyclyl, (heterocyclyl)alkyl, or (heteroaryl)alkyl);

WO 2005/051931


PCT/IB2004/003893

100


- 15 aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl) alkyl; (heterocyclyl) alkyl; $(CH_2)_{1-4}OR'$
 16 (wherein R' is as defined above, but also including hydroxy); $C(=O)NR_xR_y$
 17 (wherein R_x and R_y can be independently selected from hydrogen, alkyl, C_{3-6}
 18 alkenyl, C_{3-6} alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heteroaryl,
 19 heterocyclyl, heteroarylalkyl, or heterocyclylalkyl); or
 20 $(CH_2)_m-C(=O)R_3$
 21 [wherein m is an integer in the range of 0-2 and R_3 can be optionally substituted
 22 R_p or R_q (wherein R_p can be a 4-12 membered (un)saturated monocyclic or
 23 bicyclic ring containing 1-4 heteroatom(s) selected from N, O and S wherein the
 24 ring can be attached to $(CH_2)_mC(=O)$ through N and R_q can be a 4-12 membered
 25 (un)saturated monocyclic or bicyclic ring containing 0-4 heteroatom(s) selected
 26 from the group consisting of N, O and S wherein the ring can be attached to
 27 $(CH_2)_mC(=O)$ through C) and wherein the substituents of R_3 can be one or more
 28 of: alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl, halogen, hydroxyl, alkoxy,
 29 aryloxy, nitro, cyano, amino, substituted amino, hydroxyalkyl, oxo, acyl,
 30 optionally substituted amino (wherein the substituents are selected from C_1-C_6
 31 alkyl, aryl, aralkyl, or cycloalkyl), aryl, carboxyl, alkaryl, carbamoyl, alkyl ether,
 32 $C(=O)NR_5R_6$ (wherein R_5 and R_6 are independently selected from hydrogen,
 33 alkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, aryl, and aralkyl), optionally substituted
 34 monocyclic or bicyclic 4-12 membered carbocyclic ring system (wherein the
 35 optional substituent(s) is/are selected from alkyl, alkenyl, alkynyl, halogen,
 36 hydroxyl, and alkoxy), heteroaryl, heterocyclyl, heteroarylalkyl, or
 37 heterocyclylalkyl];
 38 R_4 is selected from: hydrogen; alkyl; halogen; cyano; carboxy; or $C(=O)NR_xR_y$ wherein
 39 R_x and R_y are the same as defined above;
 40 X_1 and X_2 are independently selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl;
 41 acyl; aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl)alkyl; or (heterocyclyl)alkyl;
 42 Y_1 and Y_2 are independently selected from: hydrogen; alkyl; nitro; cyano; halogen; OR
 43 wherein R is the same as defined earlier; SR wherein R is the same as defined earlier;
 44 NHR wherein R is the same as defined earlier; COOR'; or COR' wherein R' is the same
 45 as defined above, or further, Y_1 and X_2 , X_1 and Y_2 , X_1 and X_2 may together form a ring

46 fused with the ring A containing 3-5 carbon atoms within the ring and having 1-3
47 heteroatoms selected from N, O or S;

48 the method comprising:
49 reacting a compound of Formula VIII




56 with ethylmethylketone to give a compound of Formula XXVIII; and



63 treating the compound of Formula XXVIII with acetic anhydride to give the
64 compound of Formula XXIX.

1 27. A process for the preparation of compounds of Formula XXX,



7 their pharmaceutically acceptable salts, pharmaceutically acceptable solvates,
8 enantiomers, diastereomers or N-oxides,
9 wherein

10 R₁ is selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl; cyano; nitro; amino;
11 substituted amino; hydroxyl; alkoxy; aryloxy; COR'; COOR'

12 (wherein R' can be hydrogen, alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl,

WO 2005/051931

PCT/IB2004/003893

102


13 aryl, aralkyl, heterocyclyl, (heterocyclyl)alkyl, or (heteroaryl)alkyl);
 14 aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl) alkyl; (heterocyclyl) alkyl; (CH₂)₁₋₄OR'
 15 (wherein R' is as defined above, but also including hydroxy); C(=O)NR_xR_y
 16 (wherein R_x and R_y can be independently selected from hydrogen, alkyl, C₃₋₆
 17 alkenyl, C₃₋₆ alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heteroaryl,
 18 heterocyclyl, heteroarylalkyl, or heterocyclylalkyl); or
 19 (CH₂)_m-C(=O)R₃
 20 [wherein m is an integer in the range of 0-2 and R₃ can be optionally substituted
 21 R_p or R_q (wherein R_p can be a 4-12 membered (un)saturated monocyclic or
 22 bicyclic ring containing 1-4 heteroatom(s) selected from N, O and S wherein the
 23 ring can be attached to (CH₂)_mC(=O) through N and R_q can be a 4-12 membered
 24 (un)saturated monocyclic or bicyclic ring containing 0-4 heteroatom(s) selected
 25 from the group consisting of N, O and S wherein the ring can be attached to
 26 (CH₂)_mC(=O) through C) and wherein the substituents of R₃ can be one or more
 27 of: alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl, halogen, hydroxyl, alkoxy,
 28 aryloxy, nitro, cyano, amino, substituted amino, hydroxyalkyl, oxo, acyl,
 29 optionally substituted amino (wherein the substituents are selected from C₁-C₆
 30 alkyl, aryl, aralkyl, or cycloalkyl), aryl, carboxyl, alkaryl, carbamoyl, alkyl ether,
 31 C(=O)NR₅R₆ (wherein R₅ and R₆ are independently selected from hydrogen,
 32 alkyl, C₃₋₆ alkenyl, C₃₋₆ alkynyl, aryl, and aralkyl), optionally substituted
 33 monocyclic or bicyclic 4-12 membered carbocyclic ring system (wherein the
 34 optional substituent(s) is/are selected from alkyl, alkenyl, alkynyl, halogen,
 35 hydroxyl, and alkoxy), heteroaryl, heterocyclyl, heteroarylalkyl, or
 36 heterocyclylalkyl];
 37 R₄ is selected from: hydrogen; alkyl; halogen; cyano; carboxy; or C(=O)NR_xR_y wherein
 38 R_x and R_y are the same as defined above;
 39 X₁ and X₂ are independently selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl;
 40 acyl; aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl)alkyl; or (heterocyclyl)alkyl;
 41 Y₁ and Y₂ are independently selected from: hydrogen; alkyl; nitro; cyano; halogen; OR
 42 wherein R is the same as defined earlier; SR wherein R is the same as defined earlier;
 43 NHR wherein R is the same as defined earlier; COOR'; or COR' wherein R' is the same

WO 2005/051931

PCT/IB2004/003893


103

- 44 as defined above, or further, Y₁ and X₂, X₁ and Y₂, X₁ and X₂ may together form a ring
 45 fused with the ring A containing 3-5 carbon atoms within the ring and having 1-3
 46 heteroatoms selected from N, O or S;
 47 the method comprising reacting a compound of Formula VIII



Formula VIII

- 48
 49 with carbon disulphide to give the compound of Formula XXX.
 1 28. A method for the preparation of compounds of Formula XXXI,



Formula XXXI

- 7
 8 their pharmaceutically acceptable salts, pharmaceutically acceptable solvates,
 9 enantiomers, diastereomers or N-oxides,
 10 wherein
 11 R₁ is selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl; cyano; nitro; amino;
 12 substituted amino; hydroxyl; alkoxy; aryloxy; COR'; COOR'
 13 (wherein R' can be hydrogen, alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl,
 14 aryl, aralkyl, heterocyclyl, (heterocyclyl)alkyl, or (heteroaryl)alkyl);
 15 aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl) alkyl; (heterocyclyl) alkyl; (CH₂)₁₋₄OR'
 16 (wherein R' is as defined above, but also including hydroxy); C(=O)NR_xR_y
 17 (wherein R_x and R_y can be independently selected from hydrogen, alkyl, C₃₋₆
 18 alkenyl, C₃₋₆ alkynyl, (un)saturated cycloalkyl, aryl, aralkyl, heteroaryl,
 19 heterocyclyl, heteroarylalkyl, or heterocyclylalkyl); or

WO 2005/051931

PCT/IB2004/003893

104

20 $(\text{CH}_2)_m\text{-C(=O)R}_3$

21 [wherein m is an integer in the range of 0-2 and R_3 can be optionally substituted
 22 R_p or R_q (wherein R_p can be a 4-12 membered (un)saturated monocyclic or
 23 bicyclic ring containing 1-4 heteroatom(s) selected from N, O and S wherein the
 24 ring can be attached to $(\text{CH}_2)_m\text{C(=O)}$ through N and R_q can be a 4-12 membered
 25 (un)saturated monocyclic or bicyclic ring containing 0-4 heteroatom(s) selected
 26 from the group consisting of N, O and S wherein the ring can be attached to
 27 $(\text{CH}_2)_m\text{C(=O)}$ through C) and wherein the substituents of R_3 can be one or more
 28 of: alkyl, alkenyl, alkynyl, (un)saturated cycloalkyl, halogen, hydroxyl, alkoxy,
 29 aryloxy, nitro, cyano, amino, substituted amino, hydroxyalkyl, oxo, acyl,
 30 optionally substituted amino (wherein the substituents are selected from $\text{C}_1\text{-C}_6$
 31 alkyl, aryl, aralkyl, or cycloalkyl), aryl, carboxyl, alkaryl, carbamoyl, alkyl ether,
 32 $\text{C(=O)NR}_5\text{R}_6$ (wherein R_5 and R_6 are independently selected from hydrogen,
 33 alkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, aryl, and aralkyl), optionally substituted
 34 monocyclic or bicyclic 4-12 membered carbocyclic ring system (wherein the
 35 optional substituent(s) is/are selected from alkyl, alkenyl, alkynyl, halogen,
 36 hydroxyl, and alkoxy), heteroaryl, heterocyclyl, heteroarylalkyl, or
 37 heterocyclylalkyl];

38 R_4 is selected from: hydrogen; alkyl; halogen; cyano; carboxy; or $\text{C(=O)NR}_x\text{R}_y$, wherein
 39 R_x and R_y are the same as defined above;

40 X_1 and X_2 are independently selected from: hydrogen; alkyl; alkenyl; alkynyl; cycloalkyl;
 41 acyl; aryl; aralkyl; heteroaryl; heterocyclyl; (heteroaryl)alkyl; or (heterocyclyl)alkyl;

42 Y_1 and Y_2 are independently selected from: hydrogen; alkyl; nitro; cyano; halogen; OR
 43 wherein R is the same as defined earlier; SR wherein R is the same as defined earlier;
 44 NHR wherein R is the same as defined earlier; COOR' ; or COR' wherein R' is the same
 45 as defined above, or further, Y_1 and X_2 , X_1 and Y_2 , X_1 and X_2 may together form a ring
 46 fused with the ring A containing 3-5 carbon atoms within the ring and having 1-3
 47 heteroatoms selected from N, O or S;

48 the method comprising:

WO 2005/051931

PCT/TB2004/003893

105

49 treating a compound of Formula XXX

50


51

52

53

54

55



Formula XXX

56 with hydrazine hydrate to give the compounds of Formula XXXI.